

Cupra Super

Reg. No. N-AR 1823 Act No. 36 of 1947

Reg. No. L9906 Act No. 36 of 1947

FRAC FUNGICIDE GROUP CODE M1

A copper-containing suspension concentrate fungicide and bactericide for the control of diseases in crops as indicated



Active Ingredient:

Copper hydroxide
(equivalent to 120 g/ℓ metallic copper)

180 g/ℓ

Product approved for use in organic farming according to EEC889 and NOP by CERES GmbH



InteliGro
intelligent crop solutions

Distributed by:
Inteligro (Pty) Ltd.,
3 Marchand Street, Wellington, 7655
PO Box 85, Wellington, 7654
Tel: 021 873 6177

VERSION ROLFES AGR: 07/2018

UN NO.: 3082
SEE CONTAINER
FOR BATCH DETAILS
COUNTRY OF ORIGIN:
REPUBLIC OF SOUTH AFRICA

Rolfes 
Agri

Registered and Manufactured by:
Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07)
P.O.Box 589, Silverton, 0127, Gauteng
Tel: (012) 803-0145 • Fax: (012) 803-8418
www.rolfesagri.co.za



HARMFUL



LIFT HERE

SHAKE WELL Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use

WARNINGS:

Withholding periods: Minimum number of days between last application and harvest:

- | | |
|------------------|-------------|
| • Apples | 14 days |
| • Avocados | 14 days |
| • Beans | 3 days |
| • Citrus | 14 days |
| • Coffee | 14 days |
| • Cruciferae | 3 days |
| • Grapes (table) | 4 - 6 weeks |
| • Grapes (wine) | 21 days |
| • Mangoes | 14 days |
| • Onions | 14 days |
| • Peaches | 14 days |
| • Potatoes | 14 days |
| • Tomatoes | 3 days |
- Harmful if swallowed or inhaled.
 - Irritating to eyes, skin and respiratory tract.
 - Toxic to fish and wildlife.
 - Keep out of reach of children, uninformed persons and animals.
 - Store in a cool, dry place away from food and feed.
 - In case of poisoning, call a doctor and show him/her this label.
 - Re-entry interval: Do not enter treated area within 1 day after treatment unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or the lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Do not eat, drink or smoke whilst handling this product.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Wear rubber gloves and face shield when handling the concentrate.
- If you get it in your eyes, flush the eyes at once using clean water. If you get it on your skin, wash it off at once. If you spill it on your clothes, change and wash them.
- Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinse water to the contents of the spray tank before destroying the container.
- While spraying, avoid contact with the spray as much as you can.
- Avoid spray drift onto other crops, grazing, rivers or dams.
- Clean applicator thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams.
- Destroy the empty container by means of perforation and flattening and never re-use it for any other purpose.
- Change and wash your work clothes. Wash yourself.

RESTRICTIONS:

Max levels of copper per ha are not to be exceeded.

SYMPTOMS OF HUMAN POISONING:

Ingestion can cause diarrhoea and stomach pain. Certain individuals may be sensitive to copper. If product gets into the eyes, it causes substantial but temporary eye irritation. Inhalation may cause respiratory tract irritation.

FIRST AID TREATMENT:

Ingestion: If person is fully conscious, give large amounts of milk or water and call a doctor.

Skin contact: Wash thoroughly with soap and water.

Eye contact: Immediately flush the eyes with plenty of clean water for at least 15 minutes. Get medical attention.

Inhalation: Remove person to fresh air. Give oxygen if breathing is difficult.

NOTE TO PHYSICIAN:

Treat symptomatically.

RESISTANCE WARNING:

For resistance management, **CUPRA SUPER** is a group code M1 fungicide. Any fungus population may contain individuals naturally resistant to **CUPRA SUPER** and other group code M1 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly. These resistant fungi may not be controlled by **CUPRA SUPER** or any other group code M1 fungicide.

To delay fungicide resistance:

- Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with registered products from different fungicide group codes.
- Integrate other control methods (chemical, cultural, biological) into disease control programmes.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: USE ONLY AS INDICATED

COMPATIBILITY: Do not apply **CUPRA SUPER** in combination with any acidifying products such as zinc nitrate (ZnNO₃), ammonium sulphate (NH₄SO₄), amino-acid chelated foliar feeds or any other acidifying or chelated products, or apply these products to the fruit and leaves within 30 days after a copper application. Do not apply citric acid containing products such as potassium citrate (**CARBOLGY**[®] K) or calcium citrate in combination with or within 30 days after a **CUPRA SUPER** application.

IMPORTANT: It is essential that **CUPRA SUPER** be kept agitated throughout the entire mixing and spraying operation. **CUPRA SUPER** is a preventive spray and application must be carried out before or just as the first signs of the disease are noticed.

ENVIRONMENTAL CONDITIONS: Do not apply **CUPRA SUPER** under adverse weather conditions causing stress to the crop, such as high or low temperatures, high or low humidity, rain or frost. Do not apply when the fruit is still wet as a result of rain or dew, or if rain is imminent or expected, or under ANY circumstance where slow drying conditions might prevail or be expected as under the above conditions phytotoxicity/injury/stippling may occur. Do not apply during periods of water stress.

CITRUS: Successive **CUPRA SUPER** applications, adverse environmental conditions, late applications or acidifying products can cause stippling of fruit. To avoid stipple formation, replace every second application with mancozeb. Do not apply **CUPRA SUPER** with any buffers or acidifiers, or any other adjuvants before consulting a qualified agronomist.

CROP	DISEASE	DOSAGE PER 100 ℓ WATER	REMARKS
Apples	Scab (<i>Venturia inaequalis</i>)	170 mℓ	Apply two applications with a 7 day interval. Spray at green tip stage.
Avocados	Fruit spot (<i>Pseudo-cercospora purpurea</i>)	350 mℓ	Start applications early in November and repeat every 4 weeks. Apply 3 – 4 full cover applications during the season. Will also control Anthracnose if present at time of application.
Beans	Bacterial blight (<i>Common and halo</i>)	350 mℓ	Apply every 7 – 14 days depending on the disease pressure. Spray immediately after heavy rains, hail or dust storms.
Citrus	Black spot (<i>Guignardia citricarpa</i>)	350 mℓ	Start CUPRA SUPER applications before mid-October. Apply 3 more high volume sprays at 30 day intervals during the susceptible period from October to end December. During a high rainfall season, a fourth application may be necessary.
	Alternaria spot (<i>Alternaria alternata</i> pv. <i>citri</i>)	250 mℓ + 100 g MANCOZEB 800 WP	Apply as a high volume cover spray. Begin application at first flush (± middle September) and repeat every 28 days until harvest.
Coffee	Rust (<i>Hemileia vastatrix</i>)	550 mℓ	Apply as a full cover application, commencing before rust symptoms appear. Apply 4 - 5 sprays at 4 week intervals. Apply at least 1 000 ℓ spray mixture per hectare.
Cruciferae	Downy mildew (<i>Peronospora parasitica</i>)	350 mℓ	Apply every 7 – 14 days. Ensure good coverage of the whole plant.
Grapes	Downy mildew (<i>Plasmopara viticola</i>)	350 mℓ	Start spraying when shoots are approximately 10 cm long and repeat every 10 – 14 days depending on weather conditions. Ensure thorough coverage of the vines. Increase the spray volume progressively from 500 ℓ – 1 500 ℓ / ha, depending on the vine growth. NB: Do not apply after pea berry stage (4 - 6 weeks before harvest).
Mangoes	Bacterial spot (<i>Xanthomonas campestris</i>)	350 mℓ	Apply at 2 – 4 week intervals from just after fruit set until harvest. The product is especially suited for application as the last two sprays at the end of a spray programme.
Onions	Downy mildew (<i>Peronospora destructor</i>)	350 mℓ	Apply 3 – 4 applications with 10 – 14 day intervals. Apply in 300 – 500 ℓ water per hectare, depending on plant size. Start as soon as the disease is noticed, or when weather conditions are conducive to the development of the disease. Ensure complete spray coverage.
Peaches	Leaf curl (<i>Taphrina deformans</i>)	450 mℓ + 140 g AG-THIRAM 800 WP* (tank-mix)	Apply as a full cover spray at 60 % leaf drop and at bud swell. Add AQUA-WET** at a rate of 18 mℓ per 100 ℓ spray water.
Potatoes	Early blight (<i>Alternaria solani</i>)	350 mℓ	Commence application before or just when first symptoms are noticed. Apply every 7 – 10 days, depending on weather conditions favourable for the development of the disease. Depending on plant size, apply at 500 – 1 500 ℓ spray mixture per hectare. Ensure thorough coverage of both upper and lower leaf surfaces. Repeat application after more than 13 mm of rain or overhead irrigation.
	Late blight (<i>Phytophthora infestans</i>)	350 mℓ + 200 g MANCOZEB 800 WP (tank-mix)	Start preventive application before first signs of the disease, depending on conditions favourable for the development of late blight infections. Apply at 7 – 10 day intervals. If late blight is present in a field, spray intervals should be shortened to 5 days. Low volume application: 200 – 300 ℓ water per hectare. High volume application: Apply 1 500 ℓ spray mixture per hectare. Ensure full coverage of both upper and lower leaf surfaces. Repeat application after more than 13 mm rain or overhead irrigation.
Tomatoes	Early blight (<i>Alternaria solani</i>)	350 mℓ	Start application shortly after transplant when plants are 15 cm high and before symptoms appear. Apply every 7 – 10 days depending on weather conditions favourable for the development of the disease. Depending on plant size, apply at 500 – 2 000 ℓ spray mixture per hectare.
	Bacterial spot Bacterial speck	350 mℓ	Ensure that both the upper and lower leaf surfaces are treated. Repeat application after more than 13 mm rain or overhead irrigation. Apply as a preventive spray at 7 – 10 day intervals. Depending on plant size, apply at 500 – 1 500 ℓ spray mixture per hectare. Ensure thorough coverage of both the upper and lower leaf surfaces. Repeat application after more than 13 mm of rain or overhead irrigation.
	Bacterial speck (on seedlings)	350 mℓ	Spray seedlings at weekly intervals, using hollow cone nozzles.

*AG-THIRAM 800 WP REG. NO. L8175 ACT NO. 36 OF 1947, ROLFES AGRI (PTY) LTD

**AQUA-WET, REG. NO. L6204 ACT NO. 36 OF 1947, ROLFES AGRI (PTY) LTD