

# QUATRO

## QUATRO BORON

is a boron solution for use as both  
a foliar feed and fertigation

**INGREDIENT:**

Boron (B) 100 g/kg (130 g/l)

SG (@20 °C) = 1,30 ± 0,02



### FERTILISER GROUP 1

Reg. No. B3592, Act No. 36 of 1947



Distributed by:  
Inteligro (Pty) Ltd.,  
3 Marchand Street, Wellington, 7655  
PO Box 85, Wellington, 7654  
Tel: 021 873 6177



Registered and Manufactured by:  
Rolfes Agri (Pty) Ltd. (Reg. No. 1998/013411/07)  
P.O.Box 589, Silverton, 0127, Gauteng  
Tel: (012) 803-0145 • Fax: (012) 803-8418  
[www.rolfesagri.co.za](http://www.rolfesagri.co.za)



Product approved for the  
use in organic farming  
according to EEC889 by  
CERES GmbH.

May be used in organic farming only  
if deficiency of the micronutrient  
being applied is documented.

**UN NO.: NOT REGULATED**  
**SEE CONTAINER**  
**FOR BATCH DETAILS**  
**COUNTRY OF ORIGIN:**  
**REPUBLIC OF SOUTH AFRICA**

VERSION ROLFES AGR: 07/2018

**SHAKE WELL** Shake well before use • Store at room temperature  
Keep out of direct sunlight • Keep container closed when not in use



**Foliar feed:**  
Use enough water to achieve a full cover spray. **Do not exceed a concentration of 0,75 % (750 mℓ / 100 ℓ).** For sensitive crops do not exceed a working concentration of 0,15 % (150 mℓ / 100 ℓ).

**Soil Applications:**  
Apply as a uniform spray to the soil surface. If practical, leach **QUATRO BORON** into the root zone.

**DIRECTIONS FOR USE: USE ONLY AS DIRECTED**

APPLICATION	RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Foliar feeds	250 – 750 mℓ	100 – 200 mℓ	<b>Vegetable crops:</b> Apply as a full cover spray at 3 – 4 week intervals as required.
	50 – 150 mℓ	50 – 100 mℓ	<b>Ornamental crops:</b> Apply as a full cover spray. Repeat monthly as required.
	500 – 2000 mℓ	100 – 200 mℓ	<b>Large trees and orchards:</b> Apply as required before flowering starts and after the last harvest.
Fertigation	500 – 2000 mℓ	–	Apply with as much irrigation water as possible. Repeat application if required.
Soil application	5 – 15 ℓ	–	Apply evenly to soil and incorporate completely.

The rate of application is dependent of the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

Consult with your chemical distributor, consultant or manufacturer should crop specific programmes or any other information be required around the use of fertilisers.