

# QUATRO

## QUATRO CALCIUM

**a calcium containing suspension  
fertiliser for use as a foliar feed to  
maintain or correct calcium levels  
in plants.**

**INGREDIENT:**

Calcium (Ca) 242 g/kg (400 g/ℓ)

SG (@20 °C) = 1,69 ± 0,02

### FERTILISER GROUP 1

Reg. No. K7544, Act No. 36 of 1947



**Distributed by:**

Inteligro (Pty) Ltd.,  
3 Marchand Street, Wellington, 7655  
PO Box 85, Wellington, 7654  
Tel: 021 873 6177



Product approved for the  
use in organic farming  
according to EEC889 by  
CERES GmbH.

May be used in organic farming only  
if deficiency of the micronutrient  
being applied is documented.



**Registered and Manufactured by:**

Rolfes Agri (Pty) Ltd. (Reg. No. 1998/013411/07)  
P.O.Box 589, Silverton, 0127, Gauteng  
Tel: (012) 803-0145 • Fax: (012) 803-8418  
[www.rolfesagri.co.za](http://www.rolfesagri.co.za)

**UN NO.: NOT REGULATED  
SEE CONTAINER  
FOR BATCH DETAILS  
COUNTRY OF ORIGIN:  
REPUBLIC OF SOUTH AFRICA**

VERSION ROLFES AGRi: 07/2018



**SHAKE WELL** Shake well before use • Store at room temperature  
Keep out of direct sunlight • Keep container closed when not in use



- **QUATRO CALCIUM** should preferably be applied early in the morning or late afternoon.
- It is advisable to do a miscibility test prior to mixing **QUATRO CALCIUM** with other chemicals.
- Consult with your chemical distributor should crop specific programmes or any other information be required about the use of **QUATRO CALCIUM**.

#### RECOMMENDATIONS

Please ensure not to exceed the maximum rate per hectare. Do not exceed a concentration of 2 % (2 ℓ / 100 ℓ) if mixed with other chemicals in a spray tank.

#### Foliar Feeding:

Apply a minimum concentration of 0.2 % (2 ℓ / 1 000 ℓ water), but do not exceed a dosage rate of 2 % (2 ℓ / 100 ℓ). When low volume sprayers are used, increase the concentration accordingly so that a minimum of 2 ℓ of **QUATRO CALCIUM** is used per hectare.

#### Fertigation:

Apply 40 – 80 ℓ per hectare. The application should be split in 2 - 3 applications per season.

#### DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Grapes and persimmons	5 ℓ	200 – 400 mℓ	Apply from petal fall to one month before harvest with 10 – 14 day intervals. Apply as a full cover spray.
Nuts: Macadamias, Pecans	5 ℓ	200 – 400 mℓ	Apply from petal fall to one month before harvest with 10 – 14 day intervals. Apply as a full cover spray according to Tree Row Volume.
Pome Fruit (Apples and Pears)	5 ℓ	200 – 400 mℓ	Apply from petal fall to one month before harvest with 10 – 14 day intervals. Apply as a full cover spray according to Tree Row Volume.
Potatoes	2 ℓ	250 mℓ	Start application at tuber initiation. Apply 3 – 4 sprays at 10 – 14 day intervals. Apply in 500 ℓ water per hectare.
SUBTROPICAL FRUIT Avocados, Mangoes and Citrus	5 ℓ	250 mℓ	Apply from pea-size with monthly intervals in 2 000 ℓ water per hectare.
Soybeans, Groundnuts and Green Beans	2 ℓ	200 mℓ	Start application at flowering. Apply 2 sprays at 7-day intervals. Apply in 300 ℓ water per hectare.
STONE FRUIT (Plums, Peaches, Nectarines, Olives)	5 ℓ	200 – 400 mℓ	Apply from petal fall to one month before harvest with 10 – 14 day intervals. Apply as a full cover spray according to Tree Row Volume.
Strawberries (Field grown)	3 ℓ	250 mℓ	Start application early in the new season until one month before harvest. Apply 3 applications with 10 – 14 day intervals in 300 – 500 ℓ water per hectare.
Tomatoes and Peppers	2 ℓ	250 mℓ	Start application at first set. Repeat at 7 – 10 day intervals. Apply in 500 ℓ water per hectare.
VEGETABLES Cucurbits (Pumpkins, Watermel- ons, Cucumbers, etc.)	2 ℓ	250 mℓ	Start application directly after fruit set. Repeat every 7 – 10 days. Apply in 500 ℓ water per hectare.