

# QUATRO

## QUATRO ZINC

**a zinc containing fertilizer for use  
as a foliar feed to maintain or  
correct zinc levels in plants.**

**INGREDIENT:**

Zinc (Zn) 398 g/kg (693 g/ℓ)

SG (@20 °C) = 1,74 ± 0,02



**FERTILISER GROUP 2**  
Reg. No. B3532, Act No. 36 of 1947



Distributed by:  
**Inteligro (Pty) Ltd.**,  
3 Marchand Street, Wellington, 7655  
PO Box 85, Wellington, 7654  
Tel: 021 873 6177



Registered and Manufactured by:  
**Rolfes Agri (Pty) Ltd.** (Reg. No. 1998/013411/07)  
P.O.Box 589, Silverton, 0127, Gauteng  
Tel: (012) 803-0145 • Fax: (012) 803-8418  
[www.rolfesagri.co.za](http://www.rolfesagri.co.za)



Product approved for the  
use in organic farming  
according to EEC889 by  
CERES GmbH.

May be used in organic farming only  
if deficiency of the micronutrient  
being applied is documented.

**UN NO.: 3082**  
**SEE CONTAINER**  
**FOR BATCH DETAILS**  
**COUNTRY OF ORIGIN:**  
**REPUBLIC OF SOUTH AFRICA**



**SHAKE WELL** Shake well before use • Store at room temperature  
Keep out of direct sunlight • Keep container closed when not in use



- **QUATRO ZINC** should preferably be applied early in the morning or late afternoon.
- It is advisable to do a miscibility test prior to mixing **QUATRO ZINC** with other chemicals.
- Consult with your chemical distributor should crop specific programs or any other information be required about the use of **QUATRO ZINC**.
- Shake the container thoroughly before use.

**DIRECTIONS FOR USE:** USE ONLY AS DIRECTED.

<b>CROP</b>	<b>MAXIMUM RATE PER HECTARE</b>	<b>RATE PER 100 ℓ WATER</b>	<b>REMARKS</b>
<b>Grapes, Persimmons</b>	10 ℓ	50 – 200 mℓ	Start application when the flower buds are visible, and repeat 2 – 3 times from 10 cm shoot growth until onset on flowering. Apply in 500 - 1 000 ℓ water / ha.
<b>Pastures</b>	2 ℓ	100 – 200 mℓ	Apply 10 – 14 days prior to turnout in 200 ℓ water / ha. Keep animals out of treated pastures 10 days after application to allow for nutrient uptake.
<b>Pome Fruit:</b> Apples, Pears	10 ℓ	50 – 200 mℓ	Apply as a full cover spray at bud burst and repeat 2 – 3 times at 14 days intervals starting at 10 – 14 days after petal drop. Apply a last application after harvest, but before leaf fall. Avoid treatment during flowering.
<b>Row Crops</b> Cereals, Canola, Maize	2 ℓ	100 – 200 mℓ	Apply at 4 – 8 leave stage in 200 – 500 ℓ water / ha.
<b>Silage, Hay Pastures</b>	2 ℓ	100 – 200 mℓ	Apply when grass growth is sufficient to absorb the spray (± 15 cm high) in 500 ℓ water / ha.
<b>Stone Fruit:</b> Peaches, Plums, Apricots, Olives	10 ℓ	50 – 200 mℓ	Apply as a full cover spray at bud break and repeat 2 – 3 times at 14 day intervals from 100 % petal drop. Apply again after harvest, but before leaf fall.
<b>Subtropical Fruit</b> Avocados, Mangoes, Citrus	10 ℓ	50 – 200 mℓ	Start application at 2/3 shoot elongation. Repeat 2 – 3 times with 10 day intervals.
Bananas	10 ℓ	100 – 200 mℓ	Spray as required in 30 – 300 ℓ water / ha. Repeat if necessary.
<b>Vegetables</b>	2 ℓ	100 – 200 mℓ	Apply from 2 – 4 leaf stage in 50 – 500 ℓ water / ha.
Brassicæ	2 ℓ	400 mℓ	Apply at 4 – 8 leave stage in 50 – 500 ℓ water / ha.
Potatoes	2 ℓ	100 – 200 mℓ	Apply once 7 days after 100 % emergence in 200 – 500 ℓ water / ha.