



Zinc SC

Reg. No. K6280 Act No. 36 of 1947

FERTILISER GROUP 1

A zinc containing fertiliser for use as a foliar feed to maintain or correct zinc levels in plants.

Zinc (Zn) 397 g/kg (700 g/ℓ)

SG (@20 °C) = 1,76 ± 0,02



Product approved for the use in organic farming according to EEC889 by CERES GmbH.

UN NO.: 3082
SEE CONTAINER FOR BATCH DETAILS
COUNTRY OF ORIGIN: REPUBLIC OF SOUTH AFRICA
VERSION ROLFES AGRI: 06/2018

Rolfes 
Agri

Registered, Manufactured and Distributed by:
Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07)
P.O.Box 589, Silverton, 0127, Gauteng, RSA.
Tel: (012) 803-0145 • Fax: (012) 803-8418
www.rolfesagri.co.za

SHAKE WELL Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



- **ZINC SC** should preferably be applied early in the morning or late afternoon.
- It is advisable to do a miscibility test prior to mixing **ZINC SC** with other chemicals.
- Consult with your chemical distributor should crop specific programmes or any other information be required about the use of **ZINC SC**.

May be used in organic farming only if deficiency of the micronutrient being applied is documented.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Bananas	1 ℓ	300 mℓ	Spray as required in 30 – 300 ℓ water per hectare. Repeat if necessary.
Brassicacae	2 ℓ	400 mℓ	Apply at 4 – 8 leave stage in 50 – 500 ℓ water per hectare.
Cereals	1 ℓ	250 mℓ	Apply from tillering to when first node is detectable (Zadok's G.S. 20-31) in 200 – 500 ℓ water per hectare.
Grapes	1 ℓ	200 mℓ	Start application when the flower buds are visible, and again when the buds are separated, or at fruit set. Apply in 200 – 1 000 ℓ water per hectare.
Maize	1 ℓ	250 mℓ	Apply at 4 – 8 leave stage in 200 – 500 ℓ water per hectare.
Pastures	1 ℓ	200 mℓ	Apply 10 – 14 days prior to turnout in 200 ℓ water per hectare. Keep animals out of treated pastures for 10 days after application to allow for nutrient uptake.
POMEFRUIT: Apples, Pears	2 ℓ	100 – 200 mℓ	Apply as a full cover spray at bud burst and repeat 10 – 14 days later if necessary. Apply a last application after harvest, but before leaf fall. Avoid treatment during flowering.
Potatoes	1 ℓ	300 mℓ	Apply once 7 days after 100% emergence in 200 – 500 ℓ water per hectare.
Silage, Hay Pastures	1 ℓ	250 mℓ	Apply when grass growth is sufficient to absorb the spray (\pm 15 cm high) in 500 ℓ water per hectare.
STONEFRUIT: Peaches, Plums, Apricots, etc	2 ℓ	100 – 200 mℓ	Apply as a full cover spray at winter bud and repeat at pink bud. Apply again after harvest, but before leaf fall.