



SOIL
CONDITIONER

Novabac Flo

Reg. No. M19 Act No. 36 of 1947

FERTILISER GROUP 3

Beneficial rhizobacterial strains in a liquid formulation to improve plant growth and wheat yield.

ACTIVE INGREDIENTS

Novabac Flo contains three *Bacilli* isolates (T19, T29 and A07)

Above mentioned bacteria.....5x10⁸ colony forming units(CFU)/ml



Registered, Manufactured and Distributed
by:

Rolfes Agri (Pty) Ltd.

(Reg. No. 1998/013411/07)

288 Mundt Street, Waltloo, 0184, Gauteng,
RSA.

Tel: (012) 803-0145 • Fax: (012) 803-8418

www.rolfesagri.co.za

Expiry date (Shelf life): 6 months

SG (@20 °C) = 1,00

UN NO.: NOT REGULATED

SEE CONTAINER FOR BATCH DETAILS

COUNTRY OF ORIGIN: REPUBLIC OF SOUTH AFRICA

VERSION ROLFES AGR1: 10/2019

SHAKE WELL
Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

WARNINGS:

- Store in a cool place, dry place.
- Do not store near pesticides.
- Keep out of reach of children, uninformed persons and animals.
- Wash hands with soap and water after handling this product.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

- This product should be used as part of a balanced soil fertility and crop production program.
- The dosage rates in the table below are to be used as a guide only. The application rate is determined by soil fertility, as well as soil and crop type.
- Do not leave the product in the sun when not using. These are living organisms and are sensitive to sunlight.
- Do not use the product after the expiry date.

CROP	RECOMMENDED DOSAGE RATE	REMARKS
Wheat	Soil application: 8-16 ℓ/ha	Pre-mix with water. Mix thoroughly before applying through irrigation system.
	In-furrow application: 8-16 ℓ/ha	Pre-mix product with water before adding to application tank. Ensure thorough agitation. Apply in 25 -250 ℓ of water per hectare.
	Seed treatment: 4-8 ℓ/ton seed	Apply directly to seed. Can be diluted on a one to one ratio with water. Water volume is determined by product and seed type.