



FERTILISER



Copstar SC Fertiliser

Reg. No. B3487 Act No. 36 of 1947

FERTILISER GROUP 2

A liquid suspension concentrate fertiliser for use as a foliar feed to maintain or correct copper levels in plants.

Copper (Cu) 102 g/kg (120 g/l)

SG (@20 °C) = 1,18 ± 0,02



Registered, Manufactured and Distributed by:
Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07)
288 Mundt Street, Waltloo, 0184, Gauteng, RSA.
Tel: (012) 803-0145 • Fax: (012) 803-8418
www.rolfesagri.co.za

UN NO.: 3082
SEE CONTAINER FOR BATCH DETAILS
COUNTRY OF ORIGIN: REPUBLIC OF SOUTH AFRICA
VERSION ROLFES AGR: 01/2019

SHAKE WELL Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

- For optimum results **COPSTAR** should be used in a programme and in combination with other balanced fertilisers and foliar feeds.
- **COPSTAR** is compatible with most agricultural remedies. **It is however advisable to do a miscibility test prior to mixing with other chemicals.**
- **COPSTAR** should preferably be applied in the early morning or late afternoon. **Do not apply to plants that are undergoing a period of moisture or heat stress.**
- Consult with your chemical distributor should crop specific programmes or any other information be required about the use of **COPSTAR**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Grapes	5 ℓ	350 mℓ	Apply with bud swell and repeat application at 75 % petal fall. Apply in a minimum of 1 000 – 3 000 ℓ water per hectare.
Citrus	10 ℓ	350 mℓ	Apply in September, before flowering. Repeat 3 – 4 weeks later if necessary. Apply as a high volume application in 1 000 – 3 000 ℓ water per hectare.
Potatoes, Tomatoes, Beans and Cruciferae	5 ℓ	350 mℓ	Apply 1 – 2 applications early in the season, when there is enough leaves to absorb the spray. Repeat every 7 – 10 days if necessary. Depending on plant size apply in 500 – 1 500 ℓ water per hectare.
Onions	3 ℓ	600 mℓ	Apply from gholf ball size bulbs, and repeat 14 days later. Apply in a minimum of 500 ℓ water per hectare.
Other crops	3 ℓ	350 mℓ	Apply every 10 – 14 days if necessary to correct copper deficiencies.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.