

Not classifiable in terms of GHS

Molybdenum (Mo) 46 g/kg 72 g/ ℓ Boron (B) 3,3 g/kg 5,2 g/ ℓ

SG: (@20 °C) 1,25 ± 0,02



Registered and Manufactured: Rolfes Agri (Pty) Ltd. (Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184 Gauteng, RSA, • Tel: (012) 803-0145



PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- . Keen out of reach of children
- · Read carefully and follow all instructions.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler

FIRST AID TREATMENT:

- General Advice: The symptoms resulting from direct exposure to the product could appear a
 while after exposure. If there is persistent discomfort, seek medical attention.
- Eye contact: Flush eyes with large amount of flowing cold water for several minutes, until no
 evidence of chemical remains. If irritation persists, get medical help.
- Skin contact: Remove all contaminated clothing and shoes. Gently wipe off residual chemical and wash skin thoroughly with water and non-abrasive soap. If irritation persists, get medical helo.
- Inhalation: Remove the affected victim from exposure to an area with fresh air. If breathing
 has stopped, administer artificial respiration with the aid of a pocket mask equipped with a
 one-way valve or other proper respiratory medical device. If irritation persists, get medical
 help.
- Ingestion: If exposed or concerned, rinse mouth thoroughly with large amount of water and get medical help.

- For optimum results ACTICELL MoB should be used in a program and in combination with other balanced fertilizers and foliar feeds.
- · It is advisable to do a miscibility test prior to mixing with other chemicals.
- Consult with your chemical distributor should crop specific programs or any other information
 he required about the use of ACTICELL MoB
- Ensure thorough agitation of the product during filling and spraying. Always use spray tank strainers when pouring product into spray tank. The spray tank must be flushed out with clean water at the end of the day.
- ACTICELL MoB should preferably be applied in the early morning or late afternoon. Do not
 apply on wet fruit. Do not apply to plants that are undergoing a period of moisture or heat
 stress.
- Shake the container thoroughly before use.
- Use lower rates for maintenance and higher rates to correct deficiencies.
- Store in the original, well-sealed container away from sunlight. Keep away from children, uninformed people, animals and foodstuffs. Wash hands after use.

DIRECTIONS FOR HISE: HISE ONLY AS DIRECTED

CROP	RATE PER HECTARE	REMARKS
Dry Beans	500 – 750 ml + 250 g Ultra Bor*	Apply as full cover spray at early growth stages or before flowering and repeat as required.
Groundnuts	750 m ℓ + 250 g Ultra Bor*	Apply as full cover spray at early growth stages or before flowering and repeat as required.
Maize	500 — 750 mℓ + 1 kg Ultra Bor*	Apply as full cover spray at early growth stages or before flowering and repeat as required.
Soybeans	500 — 750 mℓ + 250 g Ultra Bor*	Apply as full cover spray at early growth stages oLbefore flowering and repeat as required.
Sunflower	500 – 750 m l + 1 kg Ultra Bor*	Apply as a full cover spray before the opening of buds.

Consult with your chemical distributor or the registration holder when applying any addition boron (ULTRA BOR) since the additional thereof depends on the boron content in the soil.

*ULTRA BOR Reg. No. B2958. Act 36 of 1947