

Acticell[®] Mango Boost



Reg. No. B4952 Act No. 36 of 1947

FERTILIZER GROUP 2

READ LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Acticell Mango Boost is a sulphur, calcium and boron fertilizer for use as a foliar feed to maintain or increase calcium and boron levels in plants.



DANGER

Sulphur	(S)	206 g/kg	299 g/ℓ
Calcium	(Ca)	75 g/kg	109 g/ℓ
Boron	(B)	3,6 g/kg	5,2 g/ℓ

HAZARD STATEMENTS:

Causes skin irritation.
May damage fertility or the unborn child.

PRECAUTIONARY STATEMENTS:

Read carefully and follow all instructions.
Store locked up.

Rolfes 
Agri

Registered, Manufactured and Distributed: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waitloo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Emergency contact details
Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Spill Response and Transport incidents
Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Store at room temperature • Keep out of direct sunlight
Keep container closed when not in use



LIFT HERE

PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read and follow all safety instructions before use.
- Wash hands and affected area thoroughly after handling.
- IF ON SKIN: Wash with plenty of water.
- If exposed or concerned, get medical advice.
- Specific treatment (see first aid on this label).
- If skin irritation occurs: Get medical help.
- Take off contaminated clothing and wash it before reuse.
- Store locked up.
- Wear protective gloves/protective clothing/eye protection/face protection/hearing protection.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler

RELEVANT HAZARDOUS SUBSTANCES:

Boric Acid <10%, Sulfur Tech 10 to 30%.

FIRST AID TREATMENT:

- **General Advice:** The symptoms resulting from direct exposure to the product could appear a while after exposure. If there is persistent discomfort, seek medical attention.
- **Eye contact:** Flush eyes with large amount of flowing cold water for several minutes, until no evidence of chemical remains. If irritation persists, get medical help.
- **Skin contact:** Remove all contaminated clothing and shoes. Gently wipe off residual chemical and wash skin thoroughly with water and non-abrasive soap. If irritation persists, get medical help.

- **Inhalation:** Remove the affected victim from exposure to an area with fresh air. If breathing has stopped, administer artificial respiration with the aid of a pocket mask equipped with a one-way valve or other proper respiratory medical device. If irritation persists, get medical help.
- **Ingestion:** If exposed or concerned, rinse mouth thoroughly with large amount of water and get medical help.
- For optimum results **ACTICELL MANGO BOOST** should be used in a program and in combination with other balanced fertilizers and foliar feeds.
- Consult with your chemical distributor should crop specific programs or any other information be required about the use of **ACTICELL MANGO BOOST**
- It is advisable to do a miscibility test prior to mixing with other chemicals.
- Ensure thorough agitation of the product during filling and spraying. Always use spray tank strainers when pouring product into spray tank. The spray tank must be flushed out with clean water at the end of the day.
- **ACTICELL MANGO BOOST** should preferably be applied in the early morning or late afternoon. Do not apply to wet fruit. Do not apply to plants that are undergoing a period of moisture or heat stress.
- Use lower rates as maintenance and higher rates to correct deficiencies.
- Store in the original, well-sealed container away from sunlight. Keep away from children, uninformed people, animals and foodstuffs. Wash hands after use.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Mango	3 – 6 ℓ	400 mℓ	Apply the first application at early flower stage and repeat 2 weeks later. Apply a third application with fruit set.
Avocado	3 – 6 ℓ	400 mℓ	Apply the first application at early flower stage and repeat 2 weeks later. Apply a third application with fruit set.
Litchis	3 – 6 ℓ	400 mℓ	Apply the first application at early flower stage and repeat 2 weeks later. Apply a third application with fruit set.
Coffee	3 – 6 ℓ	400 mℓ	Apply the first application at early flower stage and repeat 2 weeks later. Apply a third application with fruit set.
Papaya	3 – 6 ℓ	400 mℓ	Apply the first application at early flower stage and repeat 2 weeks later. Apply a third application with fruit set.