



## **HAZARD STATEMENTS:**

May damage fertility or the unborn child.

#### PRECAUTIONARY STATEMENTS:

Wear protective gloves/protective clothing/eye protection/ face protection.

IF exposed or concerned, get medical advice. Store locked up.

# **ACTIVE INGREDIENT**

Potassium	(K)	94 g/kg	131 g/ ℓ
Phosphorous	(P)	75 g/kg	105 g/l
Boron	(B)	2,2 g/kg	3 g/l
Iron	(Fe)	1,5kg	2 g/ l
Zinc	(Zn)	690 mg/kg	996 mg/{
Manganese	(Mg)	442 mg/kg	619 mg/ l
Copper	(Cu)	25 mg/kg	35 mg/ℓ
Molybdenum	(Mo)	16 mg/kg	22,4 mg/

SG: (@20 °C) 1,40 ± 0,02



Registered and Manufactured: Rolfes Agri (Pty) Ltd. (Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184 Gauteng, RSA. • Tel: (012) 803-0145

Spill Response and Transport incidents Spill Tech, Oil and chemical pollution control Tel: +27 (86) 100 0366 / +27 (83) 253 6618 www.spilltech.co.za



## PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read, and follow all safety instructions before use.
- Wear protective gloves/protective clothing/eye protection/face protection.
- IF exposed or concerned, get medical advice.
- · Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler

## RELEVANT SUBSTANCES:

Boric Acid < 10%

#### DIRECTIONS FOR IISE: LISE ONLY AS DIRECTED.

- For optimum results ACTICELL NUT: KERNEL should be used in a programme together with other products in the ACTICELL range.
- Consult with your chemical distributor, consultant or the registration holder should crop specific programmes or any other information be required about the use of ACTICELL NUT: KERNEI
- Ensure thorough agitation of the product during filling and spraying. Always use spray tank strainers when pouring product into spray tank. The spray tank must be flushed out with clean water at the end of the day.
- It is advisable to do a miscibility test prior to mixing with other chemicals.
- ACTICELL NUT: KERNEL should preferably be applied in the early morning or late afternoon.
  Do not apply to wet fruit. Do not apply to plants that are undergoing a period of moisture or heat stress.
- · Ensure good agitation throughout application.
- · Shake the container thoroughly before use.
- Use lower rates for maintenance and higher rates to correct deficiencies.
- Store in the original, well-sealed container away from sunlight. Keep away from children, uninformed people, animals, and foodstuffs. Wash hands after use.
- · Destroy empty container after use. Do not re-use for any other purpose.

CROP	RATE PER Hectare	RATE PER 100 ℓ WATER	REMARKS
Macadamias and Nut Crops	2 - 6 &	100 - 200 mℓ	Start spraying at fruit set and repeat every 2 weeks until fruit physiologically ripe. Apply in 2000 - 3000 ℓ water per hectare.