



# CARBology® AnB

Reg. No. N-F 1001, Act No. 36 of 1947

Reg. No. K8588 Act No. 36 of 1947

## FERTILISER GROUP 1

A water soluble fertilizer for use as a foliar feed on young, non-bearing trees.

Nitrogen (N)	176 g/kg	Manganese* (Mn)	9 g/kg
Phosphorus (P)	54 g/kg	Copper** (Cu)	4 g/kg
Potassium (K)	39 g/kg	Iron** (Fe)	3 g/kg
Magnesium* (Mg)	20 g/kg	Boron (B)	9 g/kg
Zinc* (Zn)	16 g/kg		

**InteliGro**  
intelligent crop solutions

**Distributed by:**  
**Inteligro (Pty) Ltd.,**  
3 Marchand Street, Wellington, 7655  
PO Box 85, Wellington, 7654  
Tel: 021 873 6177



**UN NO: NOT REGULATED**  
**SEE BLOCK ABOVE FOR BATCH DETAILS**  
**COUNTRY OF ORIGIN: REPUBLIC OF SOUTH AFRICA**  
**VERSION ROLFES AGRI: 10/2020**

**Rolfes**   
Formulating Sustainable Growth  
**Agri**

**Registered, Manufactured and Distributed by:**  
**Rolfes Agri (Pty) Ltd.**

(Reg. No. 1998/013411/07)  
288 Mundt Street, Waltloo, 0184, Gauteng, RSA.  
Tel: (012) 803-0145 • Fax: (012) 803-8418

[www.rolfesagri.co.za](http://www.rolfesagri.co.za)

**SHAKE WELL** Shake well before use • Store at room temperature  
Keep out of direct sunlight • Keep container closed when not in use



**DIRECTIONS FOR USE:** USE ONLY AS DIRECTEDDo NOT apply **CARBOLOGY® AnB**:

- In combination with copper containing fungicides, phosphates or phosphonates
  - Within 30 days after application of any copper fungicides
  - In combination with highly alkaline material such as lime sulphur, Bordeaux mixture or certain Boron products
- For use to prevent or correct nutrient deficiencies in crops as listed below.
  - Apply **CARBOLOGY® AnB** as a high volume foliar spray either in the early morning or late afternoon. Do not spray when temperatures exceed 30 °C or onto wilted plants as this can result in reduced uptake or leaf burn.
  - It is advisable to do a miscibility test prior to mixing **CARBOLOGY® AnB** with other chemicals. Do not exceed a concentration of 2 % (2 ℓ / 100 ℓ water) if mixed with other chemicals in a spray tank.
  - The use of a suitable wetter is advised.
  - **CARBOLOGY® AnB** could lower the pH of the spray water, therefore the use of a buffer is recommended when using in combination with other chemicals.

**RECOMMENDATIONS:**

CROP	RATE PER 100 ℓ WATER OR HECTARE	APPLICATION DETAILS
Grain & Row Crops	3 – 5 kg / ha	Start application at 2- 3 leaf stage (1 – 3 applications).
Nursery Tree	1 kg + suitable wetter according to label recommendations	Apply 4-6 applications, or until sufficient growth has been obtained
Pastures	3 – 5 kg / ha	Apply after every grazing or cutting
Young grapes (wine and table)	1 kg + suitable wetter according to label recommendations	Start application when sufficient leaves are present. Repeat application every 14 days, until sufficient growth has been obtained.
Young non-bearing fruit and nut trees	1 kg + suitable wetter according to label recommendations	<b>Deciduous trees:</b> Start application when sufficient leaves are present. Repeat application every 14 days.  <b>Non-deciduous trees:</b> Start application 14 days after transplant. Repeat application every 14 days.