


Cal PRO Nutech



Reg. No. B4412 Act No. 36 of 1947

Reg. No. N-F 1531 Act No. 36 of 1947

FERTILIZER GROUP 2

**READ LABEL BEFORE USE.
KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

CAL PRO NUTECH is an organic calcium fertilizer for use as a foliar feed to maintain or correct calcium levels in plants.



WARNING

Calcium* (Ca)	63 g/kg	(83 g/ℓ)
Amino acids	154 g/kg	(203 g/ℓ)
Nitrogen derived from amino acids	29 g/kg	(38 g/ℓ)
Auxin	3,7 g/kg	(4,9 g/ℓ)

*Di-amino acid chelated
SG: 1.32 ± 0.02

HAZARD STATEMENTS:

Harmful if swallowed.
Causes skin irritation.
May cause an allergic skin reaction.
Causes serious eye irritation.

PRECAUTIONARY STATEMENTS:

Read carefully and follow all instructions.
Do not eat, drink or smoke when using this product.

Rolfes 
Agri

Registered, Manufactured and Distributed by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Walthioo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Wash hands and affected area thoroughly after handling. Do **NOT** touch eyes.
- Do not eat, drink, or smoke when using this product.
- Wear eye protection/face protection.
- IF SWALLOWED: Get medical help.
- IF ON SKIN: Wash with plenty of water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- Specific treatment (see first aid treatment on this label)
- Rinse mouth.
- If eye irritation persists: Get medical help.
- Take off contaminated clothing and wash it before reuse.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

RELEVANT HAZARDOUS SUBSTANCES:

Acetic Acid < 10%, Calcium Nitrate 30 to 60%, Auxins < 10%.

FIRST AID TREATMENT:

- **General Advice:** The symptoms resulting from direct exposure to the product could appear a while after exposure. If there is persistent discomfort, seek medical attention.
- **Eye contact:** Flush eyes immediately with large amounts of flowing cold water for 15-20 minutes,

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	RATE / HA	REMARKS
Apples	6 – 12 £	Apply as a full cover spray in 1 000 – 3 000 £ spray water directly after fruit set. Repeat every 7 – 10 days to ensure 6 – 8 sprays and 6 – 12 sprays for bitter pit sensitive cultivars. If vigorous shoot growth appears use the higher dosage and increase the amount of sprays.
Avocados	6 – 12 £	Start first application on flower trusses. Repeat 3 times in early spring on new growth. Add 6 m£ AQUA-WET* per 100 £. Use the lower dosage with water volumes up to 2000 £ / ha, and the higher dosage with water volumes higher than 2000 £ / ha. Do not exceed the maximum dosage.
Brassicae, Lettuce, Beetroot	3 – 5 £	Apply 2 – 3 sprays, starting 4 weeks after transplant. Add AQUA-WET* at 6 m£ per 100 £.
Carrots	4 – 6 £	Apply as a medium cover spray and repeat at 14 day intervals.
Celery	3 – 5 £	Start application after seedlings have established. Repeat every 7 – 10 days. Add AQUA-WET* at 6 m£ per 100 £.
Cucurbits, Tomatoes, Peppers	4 – 8 £	Start application after fruit set. Repeat every 7 – 10 days. Add AQUA-WET* at 6 m£ per 100 £.
Citrus	6 – 12 £	Start application in October directly after fruit set. Repeat every 7 – 10 days (4 to 5 applications). Use the lower dosage with water volumes up to 2000 £ / ha, and the higher dosage with water volumes higher than 2000 £ / ha. Do not exceed the maximum dosage.
Cotton	3 – 6 £	Start application when plants reach a height of 30 cm. Apply 2 – 3 sprays. Add AQUA-WET* at 6 m£ per 100 £.
Grapes	4 – 6 £	Start application directly after fruit set. Apply 3 – 4 sprays. Apply in 1 000 – 1 500 £ / ha water.
Litchis, Mangoes	6 – 12 £	Start application during flowering. Apply 2 – 3 sprays at 10 day intervals. Add AQUA-WET* at 6 m£ per 100 £. Use the lower dosage with water volumes up to 2000 £ / ha, and the higher dosage with water volumes higher than 2000 £ / ha. Do not exceed the maximum dosage.
Plums, Peaches, Nectarines	6 – 12 £	Start application directly after fruit set. Apply 4 – 6 sprays at 10 day intervals. Add AQUA-WET* at 6 m£ per 100 £. Use the lower dosage with water volumes up to 1000 £ / ha, and the higher dosage with water volumes higher than 1000 £ / ha. Do not exceed the maximum dosage.
Pomegranates, Figs, Olives	6 – 12 £	Apply as a full cover spray. Start application during flowering. Repeat every 2 weeks until 2 weeks before harvesting.
Potatoes	4 – 6 £	Start application at tuber initiation. Apply 3 – 4 sprays at 10 to 14 day intervals. Add AQUA-WET* at 6 m£ per 100 £.
Soybeans, Groundnuts, Green Beans	4 – 6 £	Start application at flowering. Apply 2 sprays at 7 day intervals. Add AQUA-WET* at 6 m£ per 100 £.
Sugarcane	4 – 6 £	Start application in early spring. Apply 2 sprays at 10 day intervals. Add AQUA-WET* at 6 m£ per 100 £.

- until no evidence of chemical remains. If irritation persists, get medical help.
- **Skin contact:** Remove all contaminated clothing and shoes. Gently wipe off residual chemical and wash skin thoroughly with non-abrasive soap. If irritation persists, get medical help.
- **Inhalation:** Remove the affected victim from exposure to an area with fresh air. If breathing has stopped, administer artificial respiration with the aid of a pocket mask equipped with a one-way valve or other proper respiratory medical device.
- **Ingestion:** Seek medical attention or call a poison control centre for treatment advice. Do not induce vomiting unless instructed to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If the person is alert, rinse mouth thoroughly with water.

Do NOT apply CAL PRO NuTech:

- In combination with copper containing fungicides, phosphates or phosphonates.
- Within 30 days after application of any copper fungicides.
- In combination with highly alkaline material such as lime sulphur, MCPA or highly alkaline boron products.

CAL PRO NUTECH should preferably be applied in the early morning or late afternoon. Do not apply on wet fruit. Application can only take place once the fruit is dry. Do not apply to plants that are undergoing a period of moisture or heat stress, or during cool wet conditions.

For optimum results, CAL PRO NuTech should be used in a program and in combination with other balanced fertilizers and foliar feeds. Contact your chemical distributor or a **Rolfes Agri (Pty) Ltd** representative should crop specific programs be required, or if any other guidance is required concerning the use of this product.