

Combi PRO Nutech



Reg. No. B4406 Act No. 36 of 1947

Reg. No. N-F 1530 Act No. 36 of 1947

FERTILIZER GROUP 2

**READ LABEL BEFORE USE.
KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

COMBI PRO NUTECH is an organic fertilizer for use as a foliar feed to maintain or correct zinc, iron, copper, manganese, boron and molybdenum levels in plants.



DANGER

Zinc*	(Zn)	19 g/kg	(25 g/ℓ)
Iron*	(Fe)	29 g/kg	(38 g/ℓ)
Copper*	(Cu)	10 g/kg	(13 g/ℓ)
Manganese*	(Mn)	17 g/kg	(22 g/ℓ)
Boron	(B)	4 g/kg	(5 g/ℓ)
Molybdenum	(Mo)	136 mg/kg	(178 mg/ℓ)
Sulphur	(S)	41 g/kg	(54 g/ℓ)
Amino acids		145 g/kg	(190 g/ℓ)
Nitrogen derived		28 g/kg	(37 g/ℓ)

*Di-amino acid chelated
SG: 1.31 ± 0.02

HAZARD STATEMENTS:

Causes skin irritation.
Causes serious eye damage.
May damage fertility or the unborn child.
May cause damage to organs through prolonged or repeated exposure .
Very toxic to aquatic life.
Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS:

Wash hands and affected area thoroughly after handling.
Do not touch eyes.
Avoid release to the environment.

Rolfes 
Agri

Emergency contact details
Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Registered and Manufactured by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Spill Response and Transport incidents
Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read, and follow all safety instructions before use.
Do not breathe dust/fume/gas/mist/vapours/spray.
- Wash hands and affected area thoroughly after handling. Do **NOT** touch eyes.
- Avoid release to the environment.
- Wear protective gloves/protective clothing/safety goggles.
- IF ON SKIN: Wash with plenty of water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- Get medical help.
- IF exposed or concerned, get medical advice.
- Get medical help if you feel unwell.
- Specific treatment (see first aid on this label)
- If skin irritation occurs: Get medical help.
- Take off contaminated clothing and wash it before reuse.
- Collect spillage.
- Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

RELEVANT HAZARDOUS SUBSTANCES:

Zinc sulphate > 10%, Copper sulphate < 10%, Manganese sulphate < 10%, Ferrous sulphate 10 to 30%, Boric acid < 10%.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Do **NOT** apply **COMBI PRO NuTech**:

- In combination with copper containing fungicides, phosphates or phosphonates.
- Within 30 days after application of any copper fungicides.
- In combination with highly alkaline material such as lime sulphur, MCPA or highly alkaline boron products.

COMBI PRO NUTECH should preferably be applied in the early morning or late afternoon. Do not apply on wet fruit. Application can only take place once the fruit is dry. Do not apply to plants that are undergoing a period of moisture or heat stress, or during cool wet conditions.

For optimum results, **COMBI PRO NuTech** should be used in a program and in combination with other balanced fertilizers and foliar feeds. Contact your chemical distributor or an **Rolfes Agri (Pty) Ltd** representative should crop specific programs be required, or if any other guidance is required concerning the use of this product.

CROP	RATE PER HA	REMARKS
BERRIES AND OTHER SMALL FRUIT Grapes, Strawberries, Blackberries, Boysenberries, Cranberries, Raspberries, Blueberries, Currants	3 – 5 ℓ	Start application directly after fruit set. Apply 3 – 4 sprays every 10 days. Apply in 1 000 – 1 500 ℓ / ha.
CEREAL GRAINS Wheat, Cereal rye, Barley, Oats, Maize, (Popcorn cultivars as well), Sorghum, Millet	1 – 2 ℓ	Apply early in the season and repeat again during vegetative growth stage.
LEGUMES Soybeans, Groundnuts, Green Beans, Peas, Lucerne	1 – 3 ℓ	Start application at flowering. Apply 2 sprays at 7 day intervals.
POME FRUIT Apples, Pears	3 – 4 ℓ	Apply as a full cover spray in 1 000 – 3 000 ℓ spray water in spring when the first leaves appear. Repeat after harvest.
Pomegranates, Figs	3 – 5 ℓ	Apply as a full cover spray. Start application in early spring. Apply 2 – 3 sprays during the growing season.
Ornamentals, Lawns	2 – 4 ℓ	Apply as a medium cover spray and repeat at 14 day intervals. Apply in 500 ℓ / ha.
Stone Fruit Plums, Peaches, Nectarines, Olives	3 – 5 ℓ	Start application directly after fruit set. Apply 2 – 3 sprays at 10 day intervals. Use the lower dosage with water volumes up to 1 000 ℓ / ha, and the higher dosage with water volumes higher than 1 000 ℓ / ha. Do not exceed the maximum dosage.
Subtropical Fruit Avocados	3 – 5 ℓ	Start application in the spring, and repeat 2 – 3 times if necessary. Use the lower dosage with water volumes up to 2 000 ℓ / ha, and the higher dosage with water volumes higher than 2 000 ℓ / ha. Do not exceed the maximum dosage.
Citrus	3 – 6 ℓ	Start application early in spring. Repeat every 7 – 10 days (2 – 3 applications). Use the lower dosage with water volumes up to 2 000 ℓ / ha, and the higher dosage with water volumes higher than 2 000 ℓ / ha. Do not exceed the maximum dosage.
Litchis, Mangoes	3 – 5 ℓ	Start application shortly before flowering. Apply 2 – 3 sprays at 10 day intervals. Use the lower dosage with water volumes up to 2 000 ℓ / ha, and the higher dosage with water volumes higher than 2 000 ℓ / ha. Do not exceed the maximum dosage.
Sugarcane	2 – 4 ℓ	Start application in early spring. Apply 2 sprays at 10 day intervals.
Cotton	1 – 3 ℓ	Apply early in the season and repeat 3 - 4 times during season.

CROP	RATE PER HA	REMARKS
Sunflower	1 - 3 ℓ	Apply early in the season and repeat 2 – 3 times during season.
Nut Trees	3 – 5 ℓ	Start application directly after fruit set. Apply 3 – 5 sprays at 14 day intervals. Use the lower dosage with water volumes up to 1000 ℓ / ha, and the higher dosage with water volumes higher than 1000 ℓ / ha. Do not exceed the maximum dosage.
Macadamia nuts, Pistachios, Pecan nuts and Walnuts		
Vegetables		
Carrots	2 – 4 ℓ	Apply as a medium cover spray and repeat at 14 day intervals. Apply in 500 ℓ / ha.
Cruciferae, Lettuce, Beetroot, Celery	1 – 3 ℓ	Apply 2 – 3 sprays, starting 4 weeks after transplant.
Cucurbits (Pumpkins, Cucumbers etc.)	1 – 3 ℓ	Start application directly after fruit set. Repeat every 7 – 10 days if necessary.
Potatoes	2 – 4 ℓ	Start application at tuber initiation. Apply 3 – 4 sprays at 10 – 14 day intervals.
Tomatoes, Paprika, Peppers	2 – 4 ℓ	Start application at first set. Repeat at 7 – 10 day intervals.
Onions	2 – 4 ℓ	Apply as a medium cover spray and repeat at 14 day intervals. Apply in 500 ℓ / ha.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.