

Easygro Vegetative



Reg. No. K5240 Act No. 36 of 1947

READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

FERTILIZER GROUP 1

EasyGro Vegetative is a water soluble 6:1:3 (44) fertilizer with micro elements for foliar feeding and fertigation.



DANGER

Nitrogen (N)	265 g/kg		
Phosphorus (P)	44 g/kg		
Potassium (K)	135 g/kg	Cytokinins	2,0 mg/kg
Sulphur (S)	13 g/kg	Auxins	0,5 mg/kg
Magnesium (Mg)	2 g/kg		
*Zinc (Zn)	500 mg/kg		
*Iron (Fe)	500 mg/kg		
*Manganese (Mn)	200 mg/kg		
*Copper (Cu)	200 mg/kg		
Boron (B)	1000 mg/kg		
Molybdenum (Mo)	50 mg/kg		

*EDTA Chelated

HAZARD STATEMENTS:

May intensify fire; oxidizer.
May damage fertility or the unborn child.

PRECAUTIONARY STATEMENTS:

Keep away from heat, hot surfaces, sparks, open flames, and other ignition sources. No smoking.
If exposed or concerned, get medical aid.

Rolfes 
Agri

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Registered, Manufactured and Distributed by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Walthoo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Store at room temperature • Keep out of direct sunlight
Keep container closed when not in use



LIFT HERE

PRECAUTIONS

- If medical advice is needed, have container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read and follow all safety instructions before use.
- Keep away from heat, hot surfaces, sparks, open flames, and other ignition sources.
- No smoking.
- Keep away from clothing and other combustible product.
- Wear eye protection/face protection.
- In case of fire: Use suitable medium to extinguish
- If exposed or concerned, get medical aid.
- Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler. Do not use empty containers for any other purpose.

Relevant Substances:

Boric Acid < 10% and Potassium Nitrate 30 - 60%.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

EASYGRO VEGETATIVE is recommended for use during plant growth periods that require relatively high nitrogen levels such as during periods of vegetative growth or directly after harvest when nutrient reserves are being stored for the next season.

EASYGRO VEGETATIVE is a completely water-soluble fertilizer and chlorine free.

EASYGRO VEGETATIVE is readily taken up by the plant and is formulated for use as both a foliar feed and for fertigation.

For optimum results **EASYGRO VEGETATIVE** should be used in a program in combination with the other **EASYGRO** range of fertilizers.

Whenever the calcium and magnesium levels need to be increased, **EASYGRO VEGETATIVE** be used in combination with **EASYGRO CALMAG + TE***.

EASYGRO VEGETATIVE is compatible with most agrochemicals. **It is however advisable to do a miscibility test prior to mixing with other chemicals.**

APPLICATION	RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Foliar feed	1 – 3 kg	1 – 2 kg	Vegetable and ornamental crops. Apply as a full cover spray using ground rig or knapsack application. Apply at 1 -2 week intervals until the start of flowering.
	1 – 3 kg	1 – 3 kg / 30 ℓ water	Aerial Application. Apply at 1 – 2 week intervals as required until the start of flowering.
	4 – 6 kg	1 – 2 kg	Large trees and orchards. Apply at 1 – 2 week intervals during flowering and fruiting until the last harvest.
Fertigation	5 – 10 kg	0,25 – 2 kg	Use lower concentrations young plants and increase the rate as the plants mature. Use during periods that require a high nitrogen level.

The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

EASYGRO VEGETATIVE should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress. Consult with your chemical distributor, consultant or manufacturer should crop specific programs or any other information be required around the use of the **EASYGRO** range of fertilizers.