

EasyCrop

5:10:2 (26) + 3, 1 % MG + 4, 1 % S
+ MICRO ELEMENTS

Reg. No. K6483 Act No. 36 of 1947

FERTILIZER GROUP 1

READ LABEL BEFORE USE.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

EasyCrop is a water soluble fertilizer 5 : 10 : 2 (26) with micro elements for foliar feeding and fertigation.



DANGER

Nitrogen	(N)	78 g/kg
Phosphorus	(P)	149 g/kg
Potassium	(K)	31 g/kg
Sulphur	(S)	41 g/kg
Magnesium	(Mg)	31 g/kg
*Zinc	(Zn)	450 mg/kg

*Iron	(Fe)	3510 mg/kg
*Manganese	(Mn)	702 mg/kg
*Copper	(Cu)	56 mg/kg
Boron	(B)	1027 g/kg
Molybdenum	(Mo)	78 mg/kg

*EDTA Chelated

UN Number: 1477

Registered, Manufactured & Distributed by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waitloo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Rolfes
Agri

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za



Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read, and follow all safety instructions before use.
- Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking.
- Keep away from clothing and other combustible materials.
- Wear protective gloves/protective clothing/eye protection/face protection/hearing protection.
- IF exposed or concerned, get medical advice.
- In case of fire: Use appropriate media to extinguish.
- Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

RELEVANT HAZAROUS SUBSTANCES:

KNO₃<10% and BoricAcid<10%.

FIRST AID TREATMENT:

- **General Advice:** The symptoms resulting from direct exposure to the product could appear a while after exposure. If there is persistent discomfort, seek medical attention.

- **Eye contact:** Flush eyes with flowing cold water for several minutes, until no evidence of chemical remains. If irritation persists, get medical help.
- **Skin contact:** Remove all contaminated clothing and shoes. Gently wipe off residual chemical and wash skin thoroughly with water and non-abrasive soap. If irritation persists, get medical help.
- **Inhalation:** Remove the affected victim from exposure to an area with fresh air. If breathing has stopped, administer artificial respiration with the aid of a pocket mask equipped with a one-way valve or other proper respiratory medical device. If irritation persists, get medical help.
- **Ingestion:** If exposed or concerned, rinse mouth thoroughly with large amount of water and get medical help.
- **EASY CROP 5:10:2 (26)** should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.
- Consult with your chemical distributor, consultant or manufacturer should crop specific programs or any other information be required around the use of **EASY CROP 5:10:2 (26)**.
- **EASY CROP 5: 10:2 (26)** is a completely water-soluble fertilizer and chlorine free.
- **EASY CROP 5:10:2 (26)** is readily taken up by the plant and is formulated for use as both a foliar feed and for fertigation.
- **EASY CROP 5:10:2 (26)** is compatible with most agrochemicals. **It is however advisable to do a miscibility test prior to mixing with other chemicals.**

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

CROP	APPLICATION RATE	REMARKS
Citrus	500 g / 100 ℓ water	Apply during periods of stress.
Dry beans	4 kg / ha	Start application at the 8 leaf stage. Repeat every 14 days for a total of 3 applications.
Maize	4 kg / ha	Start application 4 – 6 weeks after emergence.
Mangoes	500 g / 100 ℓ water	Apply 2 weeks after bud break and repeat 14 days later.
Potatoes	4 kg / ha	Apply after 80% emergence. Repeat every 14 days for a total of 5 applications.
Soya beans	4 kg / ha	First application at 8 leaf stage and repeat 14 days later.
Sugarcane	4 kg / ha	Apply 12 – 16 weeks after harvest.
Sunflowers	5 kg / ha	Apply 4 – 6 weeks after emergence and repeat 14 days later.
Tobacco	4 kg / ha	Start application 4 weeks after transplant. Repeat every 7 days for a total of 3 applications.
Vegetables	4 kg / ha	Apply after 80% emergence. Repeat every 14 days for a total of 5 applications.
Wheat	4 kg / ha	Apply 6 – 8 weeks after emergence.

The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.