



Iron\* (Fe)
Sulphur (S)
Amino acids
Nitrogen derived
from amino acids

60 g/kg 35 g/kg 156 g/kg 31g/kg (74 g/ℓ) (43 g/ℓ) (192 g/ℓ) (38 g/ℓ)

## **HAZARD STATEMENTS:**

Harmful if swallowed. Causes skin irritation. Causes serious eve irritation.

### PRECAUTIONARY STATEMENTS:

Wash hand and affected area thoroughly after handling. Do not eat, drink or smoke when using this product.

\*Amino acid chelated SG: 123 + 0.02 Rolfes Agri

Registration holder: Rolfes Agri (Pty) Ltd. (Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184 Gauteng, RSA. • Tel: (012) 803-0145

Emergency contact details Office Hour Poisoning Helpline Rolfes Agri (Pty) Ltd. Tel: +27 (12) 803 0145 Spill Response and Transport incidents Spill Tech, Oil and chemical pollution control Tel: +27 (86) 100 0366 / +27 (83) 253 6618 www.spilltech.co.za



#### PRECAUTIONS.

- Wash hands and affected area thoroughly after handling. Do NOT touch eyes.
- Do not eat, drink, or smoke when using this product.
- Wear protective gloves/protective clothing/eye protection/face protection.
- IF SWALLOWED: Get medical help.
- IF ON SKIN: Wash with plenty of water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present
  and easy to do. Continue rinsing.
- Rinse mouth.
- If skin irritation occurs: Get medical help.
- Take off contaminated clothing and wash it before reuse.
  - If eye irritation persists: Get medical help.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

### RELEVANT HAZARDOUS SUBSTANCES:

Formalin < 10%. Acetic acid < 10% and Iron (II) sulfate heptahydrate 10 - 30%.

#### Do NOT apply IRON PRO NuTech:

- In combination with copper containing fungicides, phosphates or phosphonates.
- Within 30 days after application of any copper fungicides.
- In combination with highly alkaline material such as lime sulphur, MCPA or highly alkaline boron products.

IRON PRO NuTech should preferably be applied in the early morning or late afternoon. Do not apply on wet fruit. Application can only take place once the fruit is dry. Do not apply to plants that are undergoing a period of moisture or heat stress, or during cool wet conditions.

For optimum results, IRON PRO NuTech should be used in a program and in combination with other balanced fertilizers and foliar feeds. Contact your chemical distributor or a Rolfes Agri (Pty) Ltd representative should crop specific programs be required, or if any other guidance is required concerning the use of this product.

# **DIRECTIONS FOR USE:** USE ONLY AS DIRECTED

CROP	RATE / HA	REMARKS
Citrus	3−6ℓ	Two applications with $7-14$ day intervals. Do not apply after fruit set. Use the lower dosage with water volumes up to $2\ 000\ \ell$ / ha, and the higher dosage with water volumes higher than $2\ 000\ \ell$ / ha. Do not exceed the maximum dosage.
Potatoes, Tomatoes, Peppers	1 − 3 ℓ	Apply during early growth stage when enough leaves have developed for absorption.
Grapes	1 − 3 ℓ	Apply weekly until deficiency symptoms disappear. Do not apply during flowering. Apply in 1 000 − 1 500 ℓ / ha water.
Pome and Stone Fruit	2-4 ℓ	Three applications with $7-10$ day intervals. Apply first application when first leaves appear in spring. Use the lower dosage with water volumes up to $1\ 000\ \ell$ / ha, and the higher dosage with water volumes higher than $1\ 000\ \ell$ / ha. Do not exceed the maximum dosage.
Pomegranates, Figs, Olives	1 − 3 ℓ	Apply when necessary.
Sugarcane	1 − 3 ℓ	Apply early in the season when enough leaves have developed for absorption.
Wheat, Maize, Canola	1 − 3 ℓ	Contact your Rolfes Agri (Pty) Ltd representative.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.