



# **WARNING**

# **ACTIVE INGREDIENT**

Methiocarb (carbamate) Metaldehyde 5 g/kg 10 g/kg Suspected of damaging fertility or the unborn child. Very toxic to aquatic life. Very toxic to aquatic life with long lasting effects.

## PRECAUTIONARY STATEMENTS:

Avoid release to the environment. Wear protective gloves, protective clothing and eye protection when handling this product.

Registration Holder: Rolfes Agri (Pty) Ltd. (Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184 Gauteng, RSA. • Tel: (012) 803-0145



Spill Response and Transport incidents
Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za





### MOER-TU® is a Registered Trademark of Rolfes Agri (Ptv) Ltd

## WARNINGS

- Do NOT sprinkle MOER-TU® over or onto edible crops for human or animal consumption.
   Handle with care.
- CAUTION: Dogs and other domestic animals may find this product palatable.

## DISCLAIMER:

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

#### PRFCAUTIONS:

- If medical advice is needed, have container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read, and follow all safety instructions before use.
- Wear protective gloves/ protective clothing/ eve protection when handling this product.
- Avoid release to the environment.
- IF exposed or concerned, get medical help.
- Collect spillage.
- Store locked-up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler. Do not use empty containers for any other purpose.

#### RELEVANT SUBSTANCES:

This product contains Methiocarb (carbamate) 5 g/kg and Metaldehyde 10 g/kg.

## RESISTANCE WARNING STATEMENT:

For resistance management, MOER-TU® is a group code 1A insecticide. Any insect population may contain individuals naturally resistant to MOER-TU® and other group code 1A insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by MOER-TU® or any other group code 1A insecticide.

### To delay insecticide resistance:

- Avoid exclusive repeated use of insecticides from the same insecticide group code.
   Alternate with registered products from different insecticide group codes.
  - Integrate other control methods (chemical, cultural, biological) into insect programs.

## **DIRECTIONS FOR USE:** USE ONLY AS DIRECTED

## Aerial application:

Ensure that the bait is evenly distributed over the treated area and that the loss of bait onto non-target areas is restricted. The following requirements are essential to meet these criteria:

- Application must be conducted by a registered Aerial Application Operator. The aircraft must be correctly calibrated according to the instructions of SANS 10118 (Aerial Application of Aericultural Pesticides).
- Bait volume should be at a desired rate of 20 kg per ha.
- 2.0 g bait granules per m<sup>2</sup> should be recovered at the target area.
- 4. When wind speed exceeds 20 km/h, stop distribution.
- 5. Ensure that the Aerial Spray Operator knows exactly which fields to apply onto.
- Assurance needs to be obtained from the Aerial Operator that the above requirements will be met and that relevant data will be compiled and kept in a logbook for future reference.

CROP AND PEST	PEST	DOSAGE RATE	REMARKS
Various Field Crops (Vegetables, grapes, canola, etc.)	Snails and slugs	20 kg / ha	Scatter the MOER-TU® pellets evenly by hand or by using granule applicators or fertilizer spreaders where snails, slugs, millipedes and isopods occur. Repeat if necessary. Do not apply to vegetables of which the edible portion is above ground, e.g. lettuce, cabbage, cauliflower, strawberries, etc. unless when in the seedling stage. If application is required in the mature stage, apply the pellets on the ground, between rows.

## NOTES:

- When applying the pellets by hand, wear suitable gloves.
- Ensure that application equipment has been suitably calibrated.