

QUATRO CALBORMO



Reg. No. K7543 Act No. 36 of 1974

FERTILIZER GROUP 1

READ LABEL BEFORE USE.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

QUATRO CALBORMO is a Calcium, Boron and Molybdenum containing fertilizer for use as a foliar feed to maintain or correct Calcium levels in plants.



DANGER

Calcium	(Ca)	221,0 g/kg	(349,2 g/ℓ)
Boron	(B)	3,2 g/kg	(5,2 g/ℓ)
Molybdate	(Mo)	0,3 g/kg	(0,5 g/ℓ)

SG (20 °C) = 1.58 ± 0.02

HAZARD STATEMENTS:

Harmful if inhaled.
May cause cancer.
May damage fertility or unborn child.

PRECAUTIONARY STATEMENTS:

Avoid breathing dust/fumes/mist/vapours/spray.
Wear protective gloves/protective clothing/eye protection/face protection/hearing protection.

Registered and Manufactured by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Distributed by: INTELIGRO (Pty) Ltd.
PO Box 85, Wellington, 7655
Tel: +27 (21) 873 6177

Rolfes 
Agri

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read and follow all safety instructions before use.
- Avoid breathing dust/fumes/gas/mist/vapours/spray
- Use only outdoors or in a well ventilated area.
- Wear protective gloves/protective clothing/eye protection/face protection/hearing protection.
- IF INHALED: Remove person to fresh air and keep comfortable for breathing.
- Get medical help.
- IF exposed or concerned, get medical advice.
- Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

RELEVANT HAZARDOUS SUBSTANCES:

Boric Acid <10% and Formalin <10%.

FIRST AID TREATMENT:

- **General Advice:** The symptoms resulting from direct exposure to the product could appear a while after exposure. If there is persistent discomfort, seek medical attention.
- **Eye contact:** Flush eyes immediately with large amounts of flowing cold water for several minutes, until no evidence of chemical remains. If irritation persists, get medical help.
- **Skin contact:** Remove all contaminated clothing and shoes. Gently wipe off residual chemical and wash skin thoroughly with non-abrasive soap. If irritation persists, get medical help.
- **Inhalation:** Remove the affected victim from exposure to an area with fresh air. If breathing has stopped, administer artificial respiration with the aid of a pocket mask equipped with a one-way valve or other proper respiratory medical device. If irritation persists, get medical help.
- **Ingestion:** If exposed or concerned, rinse mouth thoroughly with large amounts of water and get medical advice.

QUATRO CALBORMO should preferably be applied early in the morning or late afternoon. It is advisable to do a miscibility test prior to mixing **QUATRO CALBORMO** with other chemicals. Consult with your chemical distributor should crop specific programs or any other information be required about the use of **QUATRO CALBORMO**. Shake the container thoroughly before use.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Grapes and Persimmons	20 ℓ	200 – 400 mℓ	Apply from petal fall to one month before harvest with 10 – 14 day intervals. Apply as a full cover spray.
Nuts Macadamias, Pecans	20 ℓ	200 – 400 mℓ	Apply from petal fall to one month before harvest with 10 – 14 day intervals. Apply as a full cover spray according to Tree Row Volume.
Pome Fruits Apples, Pears	5 ℓ	400 mℓ	Apply from petal fall to one month before harvest with 10 - 14 days intervals. Apply as a full cover spray according to Tree Row Volume.
Soybeans, Groundnuts and Green Beans	2 ℓ	200 mℓ	Start application at flowering. Apply 2 sprays at 7 day intervals. Apply as a full cover spray according to Tree Row Volume.
Stone Fruit Peaches, Nectarines, Plums, Olives	20 ℓ	200 – 400 mℓ	Apply from petal fall to one month before harvest with 10 – 14 day intervals. Apply as a full cover spray according to Tree Row Volume.
Subtropical Fruits Avocados, Mangoes and Citrus	5 ℓ	250 mℓ	Apply from pea-size with monthly intervals in 2 000 ℓ / ha water.
Vegetables			
Cucurbits Pumpkins, Watermelons, Cucumbers, etc.	2 ℓ	250 mℓ	Start application 2 - 4 weeks after transplant. Repeat every 7 - 10 days. Apply in 500 mℓ / ha water.
Potatoes	2 ℓ	250 mℓ	Start application at tuber initiation. Apply 3 - 4 sprays at 10 - 14 day intervals. Apply in 500 ℓ / ha water.
Tomatoes and Peppers	2 ℓ	250 mℓ	Start application at first set. Repeat at 7 – 10 day intervals. Apply in 500 ℓ / ha water.