

QUATRO LEBRAS



Reg. No. B3691 Act No. 36 of 1947

FERTILIZER GROUP 2

**READ LABEL BEFORE USE.
KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

QUATRO LEBRAS is a suspension concentrate fertilizer for use as a foliar application between stem elongation and flowering of canola, legumes, brassicas and cucurbits.



DANGER

Calcium	(Ca)	46 g/kg	(73 g/ℓ)
Sulphur	(S)	221 g/kg	(351 g/ℓ)
Zinc	(Zn)	63 g/kg	(100 g/ℓ)
Copper	(Cu)	16 g/kg	(25 g/ℓ)
Boron	(B)	32 g/kg	(50 g/ℓ)
Molybdenum	(Mo)	1 g/kg	(2 g/ℓ)

SG (20 °C) = 1.59 ± 0.02

HAZARD STATEMENTS:

Causes skin irritation.
Causes serious eye damage.
May cause cancer.
Very toxic to aquatic life.
Toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS:

Wash hands and affected area thoroughly after handling. Do not touch eyes.
Avoid release to the environment.

Registered and Manufactured by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waitloo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Distributed by: INTELIGRO (Pty) Ltd.
PO Box 85, Wellington, 7655
Tel: +27 (21) 873 6177

Rolfes 
Agri

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read label before use.
- Obtain, read and follow all safety instructions before use.
- Wash hands and affected area thoroughly after handling. Do not touch eyes.
- Avoid release to the environment.
- Wear protective gloves/protective clothing/eye protection/face protection/hearing protection.
- IF ON SKIN: Wash with plenty of water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- IF exposed or concerned, get medical advice.
- If skin irritation persists: Get medical help.
- Take off contaminated clothing and wash it before reuse.
- Collect spillage.
- Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

RELEVANT HAZARDOUS SUBSTANCES:

Zinc Oxide <10%, Sulphur 10 - 30%, Copper Oxychloride 30 - 60%, Flosperse 3000 <10%, Supragil MNS90 <10% and Formalin < 10 %.

FIRST AID TREATMENT:

- **General Advice:** The symptoms resulting from direct exposure to the product could appear a while after exposure. If there is persistent discomfort, seek medical attention.
- **Eye contact:** Flush eyes immediately with large amounts of flowing cold water for 15 - 20 minutes, until no evidence of chemical remains. If irritation persists, get medical help.
- **Skin contact:** Remove all contaminated clothing and shoes. Gently wipe off residual chemical and wash skin thoroughly with non-abrasive soap. If irritation persists, get medical help.
- **Inhalation:** Remove the affected victim from exposure to an area with fresh air. If breathing has stopped, administer artificial respiration with the aid of a pocket mask equipped with a one-way valve or other proper respiratory medical device.
- **Ingestion:** If exposed or concerned, rinse mouth thoroughly with large amount of water and get medical advice.

QUATRO LEBRAS is recommended for use during stem growth and flowering of canola, legumes, brassicas and cucurbits.

For optimum results **QUATRO LEBRAS** should be used in a program together with other balanced fertilizers and foliar feeds.

Apply **QUATRO LEBRAS** in an adequate volume of water to ensure complete coverage of the crop.

It is advisable to do a miscibility test prior to mixing with other chemicals.

QUATRO LEBRAS should preferably be applied in the early morning or late afternoon. **Do NOT apply to plants that are undergoing a period of moisture or heat stress.**

Consult your chemical distributor should crop specific programs or any other information be required regarding the use of **QUATRO LEBRAS**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	DOSAGE RATE PER HECTARE	REMARKS
Canola and legumes	1 - 2 ℓ	Apply the first application at 4 - 6 leaf stage, and repeat between stem elongation and flowering. Do not apply after flowering. Apply in a minimum of 100 ℓ / ha water.
Brassicas	1 - 2 ℓ	Apply the first application at 4 - 6 leaf stage, and repeat between stem elongation and flowering. Do not apply after flowering. Apply in a minimum of 100 ℓ / ha water.
Cucurbits	1 - 2 ℓ	Apply the first application at 4 - 6 leaf stage, and repeat between stem elongation and flowering. Do not apply after flowering. Apply in a minimum of 100 ℓ / ha water.