

# Size Up SC



Reg. No. K9656 Act No. 36 of 1947

FERTILIZER GROUP 1

READ LABEL BEFORE USE  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Size-Up SC is a water soluble liquid suspension fertilizer for foliar feeding and fertigation.

## Not classifiable in terms of GHS

Phosphorus	(P)	139 g/kg	226 g/ℓ
Potassium	(K)	176 g/kg	287 g/ℓ

SG: 20 °C = 1.63 ± 0.02

UN Number: Not Regulated

Registered and Manufactured: Rolfes Agri (Pty) Ltd.  
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184  
Gauteng, RSA. • Tel: (012) 803-0145

**Rolfes**   
**Agri**

**Emergency contact details**  
Office Hour Poisoning Helpline  
Rolfes Agri (Pty) Ltd.  
Tel: +27 (12) 803 0145

**Spill Response and Transport incidents**  
Spill Tech, Oil and chemical pollution control  
Tel: +27 (86) 100 0366 / +27 (83) 253 6618  
[www.spilltech.co.za](http://www.spilltech.co.za)

Store at room temperature • Keep out of direct sunlight  
Keep container closed when not in use



LIFT HERE

**PRECAUTIONS:**

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- **SIZE UP SC** should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.
- Clean all equipment before and after use.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

**FIRST AID TREATMENT:**

- **General Advice:** The symptoms resulting from direct exposure to the product could appear a while after exposure. If there is persistent discomfort, seek medical attention.
- **Eye contact:** Flush eyes with flowing cold water for several minutes, until no evidence of chemical remains. If irritation persists, get medical help.
- **Skin contact:** Remove all contaminated clothing and shoes. Gently wipe off residual chemical and wash skin thoroughly with water and non-abrasive soap. If irritation persists, get medical help.
- **Inhalation:** Remove the affected victim from exposure to an area with fresh air. If breathing has stopped, administer artificial respiration with the aid of a pocket mask equipped with a one-way valve or other proper respiratory medical device. If irritation persists, get medical help.
- **Ingestion:** If exposed or concerned, rinse mouth thoroughly with large amount of water and get medical help.

**RECOMMENDATIONS:**

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
<b>Bean Crops</b> (Soybean, Groundnut and Green Beans)	2 ℓ	500 mℓ	Apply as a full cover spray.
<b>Cabbage crops, Carrots</b>	2 ℓ	500 mℓ	Apply as a full cover spray.
<b>Citrus</b>	15 ℓ	200 – 500 mℓ	Apply as a full cover spray.  <b>Fruit colour and size:</b> Apply the first application after petal fall. Apply 2 – 3 applications per season. Do not apply within 30 days before harvest. Use the lower dosage on soft citrus. Application should take place in early morning when plants are not undergoing heat stress. Apply another 1 – 2 applications postharvest.
<b>Cereals</b> (Wheat, Barley, Rice, Rye, Oats)	2 ℓ	500 mℓ	Apply as a full cover spray.
<b>Cotton</b>	2 ℓ	500 mℓ	Apply as a full cover spray.
<b>Cucurbits</b> (Pumpkins, Melons, Cucumber etc.)	2 ℓ	500 mℓ	Apply as a full cover spray.
<b>Deciduous Fruit</b> (Apples, Pears, Apricots, Peaches, Cherries, Fig, Nectarine, Plums, Pomegranates)	15 ℓ	200 – 500 mℓ	Apply 2 – 3 applications after petal fall.  <b>Fruit colour and size:</b> Apply the first application four weeks prior to harvest and apply a second application 2 weeks later. Apply another 1 – 2 applications postharvest.

**DIRECTIONS FOR USE:** Use only as directed.

- Do not exceed the appropriate application rates.
- To be used only when recommended by a suitably qualified person.
- For optimum results, **SIZE UP SC** should be used in a program and in combination with other balanced fertilizers and foliar feeds.
- Consult with your chemical distributor, consultant or the registration holder should crop specific programs or any other information be required.

**Mixing instructions:**

Always follow the label recommendations. Shake container thoroughly before use. Any dosing equipment must be carefully calibrated to ensure correct and even application. Partially fill tank with water, begin agitation. Add required amount of **SIZE UP SC**. Add other chemicals if compatible. Fill tank and agitate thoroughly.

**Compatibility:**

Read all labels carefully. It is advisable to do a miscibility test prior to mixing with other chemicals. Check with your supplier before tank mixing.

**Storage:**

Keep in a cool dry chemical store, out of reach of children and livestock. Store in a well sealed container away from sunlight. Protect from cold and heat extremes.

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Wine Grapes	15 ℓ	400 – 750 mℓ	Apply 1 – 2 applications prior to flowering; typically at variation and 7 days later. Apply another 1 – 2 applications postharvest.
Table Grapes	15 ℓ	400 – 750 mℓ	Apply 1 – 2 applications prior to flowering; typically at variation and 7 days later.  <b>Berry colour and size:</b> Apply the first application four weeks prior to harvest and apply a second application 2 weeks later. Apply another 1 – 2 applications postharvest.
Flowers and Ornamentals	2 ℓ	200 mℓ	Apply as a full cover spray.
Olives	10 ℓ	500 mℓ	Apply as a full cover spray.
Lucerne and Pasture	2 ℓ	500 mℓ	Apply as a full cover spray.
Macadamias and Nut Crops	2 ℓ	500 mℓ	Apply as a full cover spray.
Maize	2 ℓ	500 mℓ	Apply as a full cover spray.
Potatoes	2 ℓ	500 mℓ	Apply as a full cover spray.
Onions	2 ℓ	500 mℓ	Apply as a full cover spray.
Subtropical Fruit (Avocados, Litchi, Mangoes, Coffee, Papaya)	2 ℓ	500 mℓ	Apply as a full cover spray.  <b>Fruit colour and size:</b> Apply the first application after petal fall. Apply 2 – 3 applications per season. Do not apply within 30 days before harvest. Application should take place in early morning when plants are not undergoing heat stress. Apply another 1 – 2 applications postharvest.
Sugarcane	2 ℓ	500 mℓ	Apply as a full cover spray.
Tomatoes, Peppers	5 – 10 ℓ	500 mℓ	Apply during the reproductive stages of the plants. The dosage rate depends on the cultivar and adequate nutritional levels during the specific growth stage.
Turf	2 ℓ	500 mℓ	Apply as a full cover spray.
Other crops not mentioned	2 ℓ	500 mℓ	Apply as a full cover spray.