

Zinc 15% EDTA Chelate



Reg. No. B3302 Act No. 36 of 1947

FERTILIZER GROUP 2

**READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

Zinc 15% EDTA CHELATE is a water-soluble powder zinc EDTA chelate for use as a soil application or foliar spray.

Not classified as hazardous according to GHS

Zinc (Zn) 150 g/kg

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Registered, Manufactured and Distributed: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Walthoo, 0184,
Gauteng, RSA. • Tel: (012) 803-0145

Rolfes 
Agri

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Store at room temperature • Keep out of direct sunlight
Keep container closed when not in use



PRECAUTIONS:

- If medical advice is needed, have container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

FOLIAR APPLICATION: Apply in sufficient water to obtain adequate coverage of the foliage. Apply early morning or late afternoon for the best results.

SOIL APPLICATION:

Apply in sufficient water to obtain even distribution. Incorporate into soil immediately after application.

COMPATIBILITY:

ZINC 15 % EDTA CHELATE is compatible with the majority of commonly used agrochemicals.

CROP	DOSAGE PER HECTARE	REMARKS
Field Crops: Maize, Beans, Peas, Soyabeans, Ground-nuts, Wheat, Barley, Oats, Cruciferae, Onions, Tomatoes	Soil: 2 – 4 kg Foliar: 1 – 2 kg	Apply immediately pre-planting. Apply annually until the deficiency is remedied, and then every 3 years to prevent recurrence. Apply early in the growing season, as soon as the crop has sufficient leaves to absorb the spray. If possible apply before deficiencies occur. Apply in a minimum of 200 ℓ water / ha.
Pecans	Foliar: 0,5 kg	Apply as a full thorough cover spray at full leaf in the spring. Repeat twice with 14 day intervals. Apply in a minimum of 500 ℓ water / ha.
Citrus	Foliar: 1 – 2 kg	Spray the new flush of growth in spring. Repeat when necessary but avoid spraying during and prior to harvest. Apply in a minimum of 500 ℓ water / ha.
Pomefruit: Apples, Pears	Foliar: 1 kg	Spray before blossom and repeat again after harvest. Apply in a minimum of 500 ℓ water / ha.
Stonefruit: Apricots, Peaches, Plums	Foliar: 1 kg Soil: 0,5 – 1 kg/tree	Spray after harvest. Do not exceed the recommended rate or leaf scorch may occur. Apply in a minimum of 500 ℓ water / ha. Apply as a drench to the drip zone area in case of severe deficiencies. Application rate depends on tree size.
Olives	Foliar: 1 kg	Spray in spring when new growth is starting, and repeat 3 – 4 weeks later. Apply in a minimum of 500 ℓ water / ha.
Cherries	Foliar: 0,5 – 1 kg	Apply as a full cover spray just after bloom. Repeat twice with 2 – 3 week intervals. Apply in a minimum of 500 ℓ water / ha.

Consult with a qualified Agronomist or Agricultural Advisor on the necessity of using it, before applying this fertilizer