

PILULA: TRACE mix



Reg. No. B5920 Act No. 36 of 1947

Reg. No. N-F 1905 Act No. 36 of 1947

FERTILIZER GROUP 2

READ LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

PILULA: Trace Mix is a pelleted fertilizer containing Fe EDTA and EDDHA as well as other micro-nutrients.



DANGER

HAZARD STATEMENTS:

May damage fertility or the unborn child.

PRECAUTIONARY STATEMENTS:

Wear protective gloves/protective clothing/eye protection/face protection.

IF exposed or concerned, get medical advice.

Iron Total	(Fe)	21 g/kg	Copper	(Cu)	2 g/kg
Zinc*	(Zn)	20 g/kg	Molybdenum	(Mo)	1 g/kg
Boron	(B)	8 g/kg			
Manganese	(Mn)	7 g/kg			
Sulphur	(S)	8 g/kg			

Registered, Manufactured & Distributed: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Walthoo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Rolfes
Agri

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Store at room temperature • Keep out of direct sunlight
Keep container closed when not in use



LIFT HERE

PRECAUTIONS AND STORAGE:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Obtain, read and follow all safety instructions before use
- Wear protective gloves/protective clothing/eye protection/face protection.
- If exposed or concerned, get medical advice.
- Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.
- Store in a cool, dry place.
- Do **NOT** store near pesticides.
- Wash hands with soap and water after handling this product.

DIRECTIONS FOR USE: Use only as directed.

- Product can be used in combination with other fertilizers.
- Spread the pellets under the tree canopy, but not directly against the stem of the tree.
- The application rates in the table below are to be used as a guide only.
- The optimal application rate is determined by soil fertility, soil type, crop type and canopy size.
- Use leaf and soil analysis as a guideline to determine the application rate since the canopy of the tree enlarges as the tree gets older.

CROP	RECOMMENDED APPLICATION RATE
<p style="text-align: center;">PERENNIAL CROPS</p>	<p style="text-align: center;">1st year: 15 g/tree 2nd year: 25 g/tree 3rd year: 40 g/tree + 4 years: 40 - 75 g/tree</p>
<p style="text-align: center;">ROW CROPS, VEGETABLES AND ORNAMENTALS</p>	<p style="text-align: center;">Broadcast: 20 kg /ha or 2 g /m²</p> <p style="text-align: center;">In-Furrow or row application: 5-10 kg /ha or 0,5 - 1 g /m²</p>
<p style="text-align: center;">TREES: PLANTING HOLE APPLICATION (1000 Trees/ha)</p>	<p style="text-align: center;">100 - 200 g/m³</p>