

Snailspray



INSECTICIDE

Reg. No. L7894 Act No. 36 of 1947

READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Snailspray is a suspension concentrate with a contact and stomach poison action for the control of snails and slugs on agricultural crops and in gardens.

IRAC INSECTICIDES GROUP CODE 1A



WARNING

HAZARD STATEMENTS:

Harmful if swallowed.
Suspected of causing cancer.
Suspected of damaging fertility or the unborn child.
Very toxic to aquatic life.

PRECAUTIONARY STATEMENTS:

Obtain, read and follow all safety instructions before use.
Avoid release to the environment.

ACTIVE INGREDIENT

Metaldehyde
Carbaryl

120 g/kg
30 g/kg

UN Number: 3082

Registration Holder: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Rolfes
Agri

Emergency contact details
Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Spill Response and Transport incidents
Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Store at room temperature • Keep out of direct sunlight
Keep container closed when not in use



WARNINGS:

- **CEASE APPLICATION 14 DAYS PRIOR TO HARVEST ON EDIBLE CROPS.**
- Toxic to fish, bees and wildlife.
- Store away from food and feeds.
- **RE-ENTRY INTERVAL** Do not enter treated area within 1 day after treatment unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read and follow all safety instructions before use.
- Wash hands and affected area thoroughly after handling.
- Do not eat, drink or smoke when using this product.
- Avoid release to the environment.
- Wear protective gloves/protective clothing/eye protection/face protection/hearing protection.
- Collect spillage.
- IF SWALLOWED: Get medical help.
- IF exposed or concerned, get medical advice.
- Rinse mouth.
- Store locked up.
- Empty all pesticides from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.
- Keep poultry out of treated area.
- Prevent contamination of food, feed, drinking water and eating utensils.

RESISTANCE WARNING:

For resistance management, **SNAILSPRAY** is a group code 1A insecticide. Any insect population may contain individuals naturally resistant to **SNAILSPRAY** and other group code 1A insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by **SNAILSPRAY** or any other group code 1A insecticide.

To delay insecticide resistance:

- Avoid exclusive repeated use of insecticides from the same insecticide group code. Alternate or tank mix with registered products from different insecticide group codes.
- Integrate other control methods (chemical, cultural, biological) into insect programs.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: USE ONLY AS INDICATED.

MIXING INSTRUCTIONS:

- **SHAKE THE CONTAINER THOROUGHLY.**
- Add the correct amount of **SNAILSPRAY** to the measured quantity of water and stir well.
- Stir the diluted suspension occasionally during application.

APPLICATION:

- Use a watering can with a fine rose.
- Mix 60 ml **SNAILSPRAY** with 10 l water and apply to 10 m².
- Ensure thorough and even wetting of both the leaves of the plants and the soil under the plants where snails and slugs hide.
- If applied by sprayer, use 60 ml **SNAILSPRAY** per 10 m² in a convenient quantity of water.
- Coarse droplets and thorough wetting is essential.
- Apply in damp weather when snails and slugs are active.
- Do not apply before rain and repeat after heavy rain.
- With heavy infestations, repeat application after 7 - 10 days.