



BoZ Pellet

Reg. No. B5919 Act No. 36 of 1947

FERTILIZER GROUP 2

READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

BoZ Pellet is a pelleted fertilizer containing Zinc and Boron.



DANGER

ACTIVE INGREDIENT

Boron	(B)	35 g/kg
Zinc*	(Zn)	112 g/kg

HAZARD STATEMENTS:

Causes serious eye damage.
May damage fertility or the unborn child.
Very toxic to aquatic life.
Very toxic to aquatic life with long-lasting effects

PRECAUTIONARY STATEMENTS:

Read carefully and follow all instructions.
Avoid release to the environment.

Rolfes 
Agri

Emergency contact details
Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Registration Holder: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Walthoo, 0184,
Gauteng, RSA. • Tel: (012) 803-0145

Spill Response and Transport incidents
Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Store at room temperature • Keep out of direct sunlight
Keep container closed when not in use



PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read, and follow all safety instructions before use.
- Wash hands and affected area thoroughly after handling. Do not touch eyes.
- Avoid release to the environment.
- Wear protective gloves/protective clothing/eye protection/face protection.
- IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- Get medical help.
- If exposed or concerned, get medical advice.
- Collect spillage.
- Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

RELEVANT HAZARDOUS SUBSTANCES:

Boric Acid <30%, Zinc Sulphate Monohydrate <50% and Sodium Benzoate <10%

FIRST AID TREATMENT

- **General Advice:** The symptoms resulting from direct exposure to the product could appear a while after exposure. If there is persistent discomfort, seek medical attention.

- **Eye contact:** Flush eyes immediately with large amounts of flowing cold water for 15 -20 minutes, until no evidence of chemical remains. If irritation persists, get medical help.
- **Skin contact:** Remove all contaminated clothing and shoes. Gently wipe off residual chemical and wash skin thoroughly with non-abrasive soap. If irritation persists, get medical help.
- **Inhalation:** Remove the affected victim from exposure to an area with fresh air. If breathing has stopped, administer artificial respiration with the aid of a pocket mask equipped with a one-way valve or other proper respiratory medical device.
- **Ingestion:** If exposed or concerned, rinse mouth thoroughly with large amount of water and get medical advice.

DIRECTIONS FOR USE: Use only as directed.

- Product can be used in-combination with other fertilizers.
- Spread the pellets under the tree canopy, but not directly against the stem of the tree.
- Use the application rates in the table below as a guide only.
- The optimal rates are determined by soil fertility, soil type, crop type and canopy size.
- Use leaf and soil analysis as guideline to determine the application rate since the canopy of the tree enlarges as the tree gets older.
- Store in a cool, dry place.
- Do not store near pesticide.

CROP	RECOMMENDED APPLICATION RATE
SUBTROPICAL FRUIT CROPS Avocados, Litchis, Mangoes, Coffee, etc.	Non-bearing and bearing trees: 20 g/m ² per season Use leaf analysis as guideline when determining the application rate
TREE NUTS Almonds, Cashew Nuts, Chestnuts, Hazelnuts, Macadamia Nuts, Pecan Nuts, Pistachio Nuts, Walnuts, etc.	Non-bearing trees: 5 g/m ² per season Bearing trees: 10 g/m ² per season
DECIDUOUS FRUIT Apples, Apricots, Cherries, Figs, Nectarine, Olives, Peaches, Pears, Plums, Pomegranates, Quinces etc.	Non-bearing trees: 5 g/m ² per season Bearing trees: 10 g/m ² per season
CITRUS FRUIT Grapefruit, Lemons, Limes, Mandarins, Oranges, Tangelos, etc.	Non-bearing trees: 5 g/m ² per season Bearing trees: 10 g/m ² per season
BERRIES AND OTHER SMALL FRUIT Blackberries, Boysenberries, Cranberries, Raspberries, Blueberries, Currants, Strawberries, Grapes, etc.	Non-bearing trees: 5 g/m ² per season Bearing trees: 10 g/m ² per season
ROW CROPS, VEGETABLES AND ORNAMENTALS	10-20 kg/ha 1-2 g/m ²