


Liquid Boron

Reg. No. B3255 Act No. 36 of 1947

FERTILIZER GROUP 1

**READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

LIQUID BORON is a boron solution for use as both a foliar feed and fertigation.



ACTIVE INGREDIENT

Boron (B) 98 g/kg (127 g/l)

SG (@20 °C) = 1.30 ± 0.02

Product approved for use in organic farming according to Reg. (EU) 2018/848: Annex II; trace elements



Registered and Manufactured by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waltho, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Emergency contact details
Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Rolfes
Agri

Spill Response and Transport incidents
Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

PRECAUTIONS:

If medical advice is needed, have product container or label at hand.

Keep out of reach of children.

Read carefully and follow all instructions.

FOLIAR FEED:

Use enough water to achieve a full cover spray. Do not exceed a concentration of 0,75 % (750 mℓ / 100 ℓ). For sensitive crops do not exceed a working concentration of 0,15 % (150 mℓ / 100 ℓ).

SOIL APPLICATIONS:

Apply as a uniform spray to the soil surface. If practical leach boron into the root zone.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

APPLICATION	RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Foliar feeds	250 – 750 mℓ	100 – 200 mℓ	Vegetable crops. Apply as a full cover spray. Apply at 3 – 4 week intervals as required.
	50 – 150 mℓ	50 – 100 mℓ	Ornamental crops. Apply as a full cover spray. Repeat monthly as required.
	500 – 2 000 mℓ	100 – 200 mℓ	Large trees and orchards. Apply as required before flowering and after the last harvest.
Fertigation	500 – 2 000 mℓ	–	Apply with as much irrigation water as possible. Repeat application as required.
Soil application	5 – 15 ℓ	–	Apply evenly to soil and incorporate completely.

The rate of application is dependent of the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies. Consult with your chemical distributor, consultant or manufacturer should crop specific programs or any other information be required around the use of fertilisers.