



# Moly Pack

Reg. No. B3522, Act No. 36 of 1947

**FERTILIZER GROUP 2**

**MOLY PACK** is a water soluble molybdenum containing powder for foliar application or seed dressing for the prevention and correction of molybdenum deficiencies.

**Not classified as hazardous according to GHS**

## ACTIVE INGREDIENT

Molybdenum (Mo) 390 g/kg

Registered and Distributed by: Rolfes Agri (Pty) Ltd.  
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waitloo, 0184  
Gauteng, RSA. • Tel: (012) 803-0145

### Emergency contact details

Office Hour Poisoning Helpline  
Rolfes Agri (Pty) Ltd.  
Tel: +27 (12) 803 0145

**Rolfes**   
**Agri**

### Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control  
Tel: +27 (86) 100 0366 / +27 (83) 253 6618  
[www.spilltech.co.za](http://www.spilltech.co.za)

Shake well before use • Store at room temperature  
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

**PRECAUTIONS:**

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.

**DIRECTIONS FOR USE:** USE ONLY AS DIRECTED

CROP	APPLICATION RATE	REMARKS
Cucurbits	2 g / ℓ	Soak the seeds overnight in a 0,2 % solution. Plant directly afterwards.
Cruciferae	100 – 250 g / ha	Apply to the foliage of young plants. Use in sufficient volume of water.
Grass / Clover Pastures	155 g / ha	Apply as a foliar application to young regrowth after cutting.
Lucerne	125 – 250 g / ha	Apply as a seed treatment with inoculation.
Maize	50 g / 1,25 ℓ water	Apply as a seed treatment before planting. Place 100 kg seed with the <b>Moly Pack</b> solution in a drum with a watertight lid. Rotate the drum for 15 – 20 minutes.
Soybeans	75 – 125 g / ha	Apply as a foliar spray to the seedlings.
Sunflower	25 g / 25 kg seed 5 g/100 ℓ water (Max. 75 g/ha)	Uniformly apply a solution of <b>Moly Pack</b> to the seed. Apply as a full cover foliar spray to the seedlings.
Tomatoes	250 g / ha	Apply as a foliar spray or to the planting furrow before transplanting. In general, molybdenum deficiencies occur throughout the Lowveld, particularly on acid soils.

**REPEAT APPLICATION:**

The following intervals between applications are recommended, depending on the quantity of **MOLY PACK** applied.

QUANTITY OF MOLY PACK APPLIED PER HECTARE	INTERVAL PERIOD BETWEEN TREATMENTS
Less than 75 g	1 Year
75 g – 150 g	2 – 3 years
More than 150 g	4 – 5 years

Consult with a qualified Agronomist or Agricultural Advisor on the necessity of using this fertilizer, before applying it.