



# Copper 15% EDTA Chelate

Reg. No. B3301 Act No. 36 of 1947

**FERTILIZER GROUP 2**

**READ LABEL BEFORE USE  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

**COPPER 15% EDTA CHELATE** is a water soluble powder copper EDTA chelate for use as a soil application or foliar spray.



**WARNING**

**HAZARD STATEMENTS:**

Causes serious eye irritation.  
Harmful if swallowed.

**PRECAUTIONARY STATEMENTS:**

Wash affected area thoroughly after handling. Do not touch eyes.  
Do not eat, drink or smoke when using this product.

Copper (Cu) 150 g/kg

**Registered, Manufactured and Distributed by:** Rolfes Agri (Pty) Ltd.  
(Reg. No. 1998/013411/07) • 288 Mundt Street, Walitoo, 0184,  
Gauteng, RSA. • Tel: (012) 803-0145

**Emergency contact details**

Office Hour Poisoning Helpline  
Rolfes Agri (Pty) Ltd.  
Tel: +27 (12) 803 0145

**Rolfes**   
**Agri**

**Spill Response and Transport incidents**

Spill Tech, Oil and chemical pollution control  
Tel: +27 (86) 100 0366 / +27 (83) 253 6618  
[www.spilltech.co.za](http://www.spilltech.co.za)

Store at room temperature • Keep out of direct sunlight  
Keep container closed when not in use



LIFT HERE

**PRECAUTIONS:**

- If medical advice is needed, have container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Wash affected area thoroughly after handling. Do **NOT** touch eyes.
- Do **NOT** eat, drink or smoke when using this product.
- Wear safety goggles and dust mask when handling this product.
- IF SWALLOWED: Get medical help.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

- Rinse mouth.
- If eye irritation persists: Get medical help.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler. Do **NOT** use empty containers for any other purpose.

**RELEVANT SUBSTANCE:**

Cu EDTA > 60%

**DIRECTIONS FOR USE:** USE ONLY AS DIRECTED.

**FOLIAR APPLICATION:** Apply in sufficient water to obtain adequate coverage of the foliage. Apply early morning or late afternoon for the best results.

**SOIL APPLICATION:**

Apply in sufficient water to obtain even distribution. Incorporate into soil immediately after application.

**COMPATIBILITY:**

**COPPER 15 % EDTA CHELATE** is compatible with the most agriculture chemicals.

**WARNING:**

The use of a wetter or buffer is not recommended as phytotoxicity may occur.

CROP	DOSAGE PER HECTARE	REMARKS
Citrus and other fruit crops	Foliar: 0,5 kg	Apply early in the season and repeat 3 – 4 weeks later. Avoid spraying during bloom and just before harvest. Apply in a minimum of 500 ℓ water/ha.
Wheat, Barley, Oats, Maize and Onions	Soil: 0,5 – 1 kg  Foliar: 0,25 - 0,5 kg	Apply immediately pre-planting in a minimum of 200 ℓ water/ha. Apply annually until the deficiency is corrected. Thereafter apply every three years to prevent recurrence.  If possible apply before deficiencies occur and repeat 3 – 4 weeks later if necessary. Apply early in the growing season, as soon as the crop has sufficient leaves to absorb the spray. Apply in a minimum of 200 ℓ water/ha.
Conifers	Foliar: 1 kg	Apply by air, ULV or conventional sprayer to second generation trees. Apply in 200 – 500 ℓ water/ha depending on tree size.
Pasture	Re-seeding: 4 kg  Established pasture: 0,5 – 1 kg	Spray and incorporate into the soil prior to re-seeding.  Spray early in the season before spring growth starts, and again between cuts of silage or hay. <b>Keep sheep out of recently sprayed pasture.</b>

Consult with a qualified Agronomist or Agricultural Advisor on the necessity of using it, before applying this fertilizer.