

Manganese 13% EDTA Chelate



Reg. No. B3300 Act No. 36 of 1947

FERTILIZER GROUP 2

READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Manganese 13% EDTA CHELATE is a water-soluble powder manganese EDTA chelate for use as a foliar spray.

Not classified as hazardous according to GHS

Manganese (Mn) 130 g/kg

EDTA Chelated

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Registered and Distributed by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184,
Gauteng, RSA. • Tel: (012) 803-0145

Rolfes 
Agri

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Store at room temperature • Keep out of direct sunlight
Keep container closed when not in use



PRECAUTIONS:

- If medical advice is needed, have container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.

COMPATIBILITY:

MANGANESE 13 % EDTA CHELATE is compatible with most foliar applied agrochemicals.

FOLIAR APPLICATION:

- Apply in sufficient water to obtain adequate coverage of the foliage.
- Apply early morning or late afternoon for the best results.
- Manganese deficiency is most likely to occur in highly leached, very sandy, alkaline, calcareous or heavy limed soils.
- Manganese deficiency is seldom encountered in loam or heavy acid soils.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	DOSAGE PER HECTARE	REMARKS
Stone fruit: Apricots, Peaches, Plums	1 kg	Apply as a foliar spray after harvest. Apply in a minimum of 500 ℓ water/ha.
Pome Fruit: Apples, Pears	1 kg	Apply as a foliar spray before blossom and repeat after harvest. Apply in a minimum of 500 ℓ water/ha.
Citrus	1 kg	Apply as a foliar spray to new growth flush in early spring. Repeat 3 – 4 weeks later. Apply in a minimum of 500 ℓ water/ha.
Field Crops: Maize, Beans, Peas, Soya beans, Groundnuts, Wheat, Barley, Oats, Cruciferae, Onions, Tomatoes	1,2 kg	Apply early in the growing season when the crop has sufficient leaves to absorb the spray. Repeat 3 – 4 weeks later if necessary. Apply in a minimum of 200 ℓ water/ha.
Olives	1,2 kg	Apply during late spring as a foliar spray in a minimum of 500 ℓ water/ha.
Pastures	4 kg	Apply early spring and repeat annually. Apply in a minimum of 200 ℓ water/ha.

Consult with a qualified Agronomist or Agricultural Advisor on the necessity of using it, before applying this fertilizer.