

# PILULA: MICRO-S



Reg. No. B5921 Act No. 36 of 1947

## FERTILIZER GROUP 2

**READ LABEL BEFORE USE**  
**KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

PILULA MICRO-S is a pelleted fertilizer containing micro-nutrients.



## DANGER

Sulphur	(S)	29 g/kg
Zinc	(Zn)	16 g/kg
Iron	(Fe)	15 g/kg
Calcium	(Ca)	10 g/kg
Boron	(B)	8 g/kg
Copper	(Cu)	5 g/kg
Manganese	(Mn)	3 g/kg
Molybdenum	(Mo)	1 g/kg

### HAZARD STATEMENTS:

Causes skin irritation.  
Causes serious eye damage.  
May cause damage to organs through prolonged or repeated exposure.  
Toxic to aquatic life.  
Toxic to aquatic life with long lasting effects.

### PRECAUTIONARY STATEMENTS:

Avoid release to the environment.  
Wear protective gloves/eye protection/face protection.

**Rolfes**   
**Agri**

**Registration Holder:** Rolfes Agri (Pty) Ltd.  
(Reg. No. 1998/01341/07) • 288 Mundt Street, Walthoo, 0184  
Gauteng, RSA. • Tel: (012) 803-0145

### Emergency contact details

Office Hour Poisoning Helpline  
Rolfes Agri (Pty) Ltd.  
Tel: +27 (12) 803 0145

### Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control  
Tel: +27 (86) 100 0366 / +27 (83) 253 6618  
[www.spilltech.co.za](http://www.spilltech.co.za)

Store at room temperature • Keep out of direct sunlight  
Keep container closed when not in use



**PRECAUTIONS:**

- Do **NOT** breathe dust.
  - Wash hands and affected area thoroughly after handling. Do **NOT** touch eyes.
  - Avoid release to the environment.
  - Wear protective gloves/eye protection/face protection.
  - IF ON SKIN: Wash with plenty of water and non-abrasive soap.
  - IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
  - Get medical help if you feel unwell.
  - If skin irritation occurs: Get medical help.
  - Take off contaminated clothing and wash it before reuse.
  - Collect spillage.
- Empty all contents from the container by placing it upside down over the spray tank and holding

it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler (South Africa).

- Store in a cool, dry place.
- Do **NOT** store near pesticides.

**DIRECTIONS FOR USE:** Use only as directed.

- Product can be used in combination with other fertilizers.
- Spread the pellets under the tree canopy, but not directly against the stem of the tree.
- The application rates in the table below are to be used as a guide only.
- The optimal application rate is determined by soil fertility, soil type, crop type and canopy size.
- Use leaf and soil analysis as a guideline to determine the application rate since the canopy of the tree enlarges as the tree gets older.

CROP	RECOMMENDED APPLICATION RATE
<p style="text-align: center;"><b>PERENNIAL CROPS</b></p>	<p style="text-align: center;">1<sup>st</sup> year: 15 g/tree            2<sup>nd</sup> year: 25 g/tree            3<sup>rd</sup> year: 40 g/tree            + 4 years: 40 - 75 g/tree</p>
<p style="text-align: center;"><b>ROW CROPS, VEGETABLES AND ORNAMENTALS</b></p>	<p style="text-align: center;"><b>Broadcast:</b>            20 kg/ha or 2 g/m<sup>2</sup></p> <p style="text-align: center;"><b>In-Furrow or row application:</b>            5 -10 kg/ha or 0.5 - 1 g/m<sup>2</sup></p>
<p style="text-align: center;"><b>TREES: PLANTING HOLE APPLICATION (1000 Trees/ha)</b></p>	<p style="text-align: center;">100 - 200 g/m<sup>3</sup></p>