



PILULA: Cu

Reg. No. B5961 Act No. 36 of 1947

FERTILIZER GROUP 2

**READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

PILULA: Cu is a pelleted fertilizer containing Cu EDTA.



WARNING

Copper (Cu) 52 g/kg

HAZARD STATEMENTS:

Causes serious eye irritation.

PRECAUTIONARY STATEMENTS:

Wash hands and affected area thoroughly after handling.

Do not touch eyes.

Wear eye protection/face protection.

Rolfes
Agri

Registered, Manufactured & Distributed by : Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Walthoo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Store at room temperature • Keep out of direct sunlight
Keep container closed when not in use



PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Wash hands and affected are thoroughly after handling. Do not touch eyes.
- Wear eye protection/face protection.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- If eye irritation persists: Get medical help.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.
- Store in a cool, dry place.
- Do **NOT** store near pesticides.

DIRECTIONS FOR USE: Use only as directed.

- Product can be used in combination with other fertilizers.
- Spread the pellets under the tree canopy, but not directly against the stem of the tree.
- The application rates in the table below are to be used as a guide only.
- The optimal application rate is determined by soil fertility, soil type, crop type and canopy size.
- Use leaf and soil analysis as a guideline to determine the application rate since the canopy of the tree enlarges as the tree gets older.

CROP	RECOMMENDED APPLICATION RATE
<p align="center">ROW CROPS, VEGETABLES AND ORNAMENTALS</p>	<p align="center">Broadcast: 15 - 20 kg/ha or 1,5 - 2 g/m²</p> <p align="center">In-Furrow or row application: 5 - 10 kg/ha or 0,5 - 1 g/m²</p>
<p align="center">TREES: MAINTENANCE</p> <p align="center">Non-Bearing Trees</p> <p align="center">Bearing Trees</p> <p align="center">General (1000 Trees/ha)</p>	<p align="center">10 – 20 g/tree</p> <p align="center">50 – 150 g/tree</p> <p align="center">100 g/tree</p>
<p align="center">TREES: DEFICIENCIES</p> <p align="center">Non-Bearing Trees</p> <p align="center">Bearing Trees</p>	<p align="center">25 – 50 g/tree</p> <p align="center">150 – 300 g/tree</p> <p align="center">Repeat 2 months later if the lower rates have been applied</p>