

Not classified as hazardous according to GHS

Zinc

(Zn)

55 a/ka





Rolfes &





## PRECAUTIONS AND STORAGE:

- If medical advice is needed, have product container or label at hand.
- . Keep out of reach of children.
- Read carefully and follow all instructions.
- Store in a cool, dry place.
- Do NOT store near pesticides.
- Wash hands with soap and water after handling this product.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

## **DIRECTIONS FOR USE:** Use only as directed.

- Product can be used in combination with other fertilizers.
- Spread the pellets under the tree canopy, but not directly against the stem of the tree.
  - The application rates in the table below are to be used as a guide only.
  - The optimal application rate is determined by soil fertility, soil type, crop type and canopy size.
- Use leaf and soil analysis as a guideline to determine the application rate since the canopy of the tree enlarges as the tree gets older.

CROP	RECOMMENDED DOSAGE RATE
ROW CROPS, VEGETABLES AND ORNAMENTALS	<b>Broadcast:</b> $15-20\ kg/ha\ or\ 1,5-2\ g\ /m^2$ <b>In-Furrow or row application:</b> $5-10\ kg/ha\ or\ 0,5-1\ g\ /m^2$
TREES: MAINTENANCE  Non-Bearing Trees  Bearing Trees	10 — 20 g/tree 50 — 150 g/tree Use lower rates for younger trees and increase rate for older trees
General (1000 Trees/ha)	100 g/tree
<b>TREES: DEFICIENCIES</b> Non-Bearing Trees  Bearing Trees	25 – 50 g/tree 150 – 300 g/tree Repeat 2 months later if the lower rates have been applied