

Zinc SC

Reg. No. K6280 Act No. 36 of 1947

FERTILIZER GROUP 1

READ LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

ZINC SC is a zinc containing fertilizer for use as a foliar feed to maintain or correct zinc levels in plants.



WARNING

HAZARD STATEMENTS:

Causes serious eye irritation.

Very toxic to aquatic life.

Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS:

Wash hands and affected area thoroughly after handling.
Do not touch eyes.

Avoid release to the environment.

Zinc (Zn) 399 g/kg (700 g/ℓ)

SG = 1,76

Product approved for use in organic farming
according to Reg. (EU) 2018/848:
Annex II; trace elements



Rolfes 
Agri

Registered, Manufactured & Distributed by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Store at room temperature • Keep out of direct sunlight
Keep container closed when not in use



PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read label before use.
- Wash hands and affected area thoroughly after handling. Do NOT touch eyes.
- Avoid release to the environment.
- Wear eye protection/face protection.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- If eye irritation persists: Get medical help.
- Collect spillage.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

RELEVANT HAZARDOUS SUBSTANCES:

Zinc Oxide 30-60%, Acrylic Poly Sodium Salt <10%

DIRECTIONS FOR USE: USE ONLY AS DIRECTED**FIRST AID TREATMENT:**

General Advice: The symptoms resulting from direct exposure to the product could appear a while after exposure. If there is persistent discomfort, seek medical attention.

- **Eye contact:** Flush eyes immediately with large amounts of flowing cold water for 15-20 minutes, until no evidence of chemical remains. If irritation persists, get medical help.
- **Skin contact:** Remove all contaminated clothing and shoes. Gently wipe off residual chemical and wash skin thoroughly with non-abrasive soap. If irritation persists, get medical help.
- **Inhalation:** Remove the affected victim from exposure to an area with fresh air. If breathing has stopped, administer artificial respiration with the aid of a pocket mask equipped with a one-way valve or other proper respiratory medical device.
- **Ingestion:** If exposed or concerned, rinse mouth thoroughly with large amounts of water and get medical advice.
- **ZINC SC** should preferably be applied early in the morning or late afternoon.
- It is advisable to do a miscibility test prior to mixing **ZINC SC** with other chemicals.
- Consult with your chemical distributor should crop specific programs or any other information be required about the use of **ZINC SC**.

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 £ WATER	RECOMMENDATIONS
POME FRUIT: Apples, Pears	2 £	100 – 200 m£	Apply as a full cover spray at bud burst and repeat 10-14 days later if necessary. Apply a last application after harvest, but before leaf fall. Avoid treatment during flowering.
STONE FRUIT: Peaches, Plums, Apricots	2 £	100 – 200 m£	Apply as a full cover spray at winter bud and repeat at pink bud. Apply again after harvest, but before leaf fall.
Bananas	1 £	300 m£	Spray as required in 30 – 300 £ water per hectare. Repeat if necessary.
Grapes	1 £	200 m£	Start application when the flower buds are visible, and again when the buds are separated, or at fruit set. Apply in 200 – 1000 £ water per hectare.
Brassica	2 £	400 m£	Apply at 4 – 8 leaf stage in 50 – 500 £ water per hectare.
Potatoes	1 £	300 m£	Apply once 7 days after 100% emergence in 200 – 500 £ water per hectare.
Maize	1 £	250 m£	Apply at 4 – 8 leaf stage in 200 – 500 £ water per hectare.
Cereals	1 £	250 m£	Apply from tillering to when first node is detectable (Zadok's G.S. 20-31) in 200 – 500 £ water per hectare.
Silage, Hay Pastures	1 £	250 m£	Apply when grass growth is sufficient to absorb the spray (\pm 15 cm high) in 500 £ water per hectare.
Pastures	1 £	200 m£	Apply 10 – 14 days prior to turnout in 200 £ water per hectare. Keep animals out of treated pastures 10 days after application to allow for nutrient uptake.