

Cop PRO Nutech



Reg. No. B4413 Act No. 36 of 1947

Reg. No. N-F 2029 Act No. 36 of 1947

FERTILIZER GROUP 2

READ LABEL BEFORE USE.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

COP PRO NUTECH is an organic copper fertilizer for use as a foliar feed to maintain or correct copper levels in plants.



DANGER

Copper*	(Cu)	102 g/kg	(141 g/ℓ)
Sulphur	(S)	56 g/kg	(77 g/ℓ)
Amino acids		154 g/kg	(213 g/ℓ)
Nitrogen derived from amino acids		29 g/kg	(40 g/ℓ)

*Di-amino acid chelated

SG: 1.38 ± 0.02

HAZARD STATEMENTS:

Harmful if swallowed.
Causes skin irritation.
Causes serious eye damage.
Very toxic to aquatic life.
Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS:

Wash hands and affected area thoroughly after handling. Do not touch eyes.
Rinse mouth.

Registered, Manufactured and Distributed by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Walthioo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Rolfes 
Agri

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



PRECAUTIONS:

- If medical advice is needed, have product container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Wash hands and affected area thoroughly after handling. Do **NOT** touch eyes.
- Do not eat, drink, or smoke when using this product.
- Avoid release to the environment.
- Wear protective gloves/eye protection/face protection.
- IF SWALLOWED: Get medical help.
- IF ON SKIN: Wash with plenty of water.
- IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- Get medical help.
- Specific treatment (see first aid on this label)
- Rinse mouth.
- If skin irritation occurs: Get medical help.
- Take off contaminated clothing and wash it before reuse.
- Collect spillage.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler.

RELEVANT HAZARDOUS SUBSTANCES:

Copper Sulphate Penta Hydrate 30 to 60% and Ammonia < 10%.

Do NOT apply COP PRO NuTech:

- In combination with copper containing fungicides, phosphates or phosphonates.
- Within 30 days after application of any copper fungicides.
- In combination with highly alkaline material such as lime sulphur, MCPA or highly alkaline boron products.

COP PRO NUTECH should preferably be applied in the early morning or late afternoon. Do not apply on wet fruit. Application can only take place once the fruit is dry. Do not apply to plants that are undergoing a period of moisture or heat stress, or during cool wet conditions.

For optimum results, **COP PRO NuTech** should be used in a program and in combination with other balanced fertilizers and foliar feeds. Contact your chemical distributor or a **Rolfes Agri (Pty) Ltd** representative should crop specific programs be required, or if any other guidance is required concerning the use of this product.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	RATE / HA	REMARKS
Citrus	500 mℓ – 1 ℓ	Apply in September, before flowering. Repeat 3 – 4 weeks later if necessary. Use the lower dosage with water volumes up to 2 000 ℓ / ha, and the higher dosage with water volumes higher than 2 000 ℓ / ha. Do not exceed the maximum dosage. Important: Use the lowest rate where copper oxychloride has been applied previously, and on non-bearing trees.
Grapes	350 – 700 mℓ	Apply as preventative and curative spray.
Onions	500 mℓ – 1 ℓ	Apply from golf ball size bulbs, and repeat 14 days later.
Pastures	200 – 750 mℓ	Apply early in the season before spring growth starts. Repeat between cuts of silage or hay. Important: Keep all livestock out of sprayed pasture for two weeks after application.
Pome, Stone, Deciduous, Subtropical Fruit	500 mℓ – 1 ℓ	Apply with bud swell and repeat application at 75% petal fall. Apply in a minimum of 1 000 – 3 000 ℓ / ha water.
Pomegranates, Figs, Olives	250 – 500 mℓ	Apply so as to address plant nutrient requirements (Cu) and to correct deficiency thereof.
Sugarbeet	200 – 750 mℓ	For best results, apply early in the season when there is sufficient leaves to absorb the spray. Apply 1 – 2 applications.
Wheat, Barley, Oats, Maize	1 – 2 ℓ	Apply 1 – 2 applications early in the season when there are enough leaves to absorb the spray.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.