

Easygro Flower & Fruit



Reg. No. K5242 Act No. 36 of 1947

FERTILIZER GROUP 1

READ LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

EasyGro Flower & Fruit a water soluble 3:1:6 (46) fertilizer with micro elements for foliar feeding and fertigation.



WARNING

HAZARD STATEMENTS:

May intensify fire; oxidizer.
May damage fertility or the unborn child.

PRECAUTIONARY STATEMENTS:

Keep away from heat, hot surfaces, sparks, open flames, and other ignition sources. No smoking.
If exposed or concerned, get medical aid.

Nitrogen (N)	141 g/kg
Phosphorus (P)	46 g/kg
Potassium (K)	275 g/kg
Sulphur (S)	13 g/kg
Magnesium (Mg)	2 g/kg
*Zinc (Zn)	500 mg/kg
*Iron (Fe)	500 mg/kg
*Manganese (Mn)	200 mg/kg
*Copper (Cu)	200 mg/kg
Boron (B)	1000 mg/kg
Molybdenum (Mo)	50 mg/kg

Cytokinins	0,5 mg/kg
Auxins	1,0 mg/kg

***EDTA Chelated
Potassium Nitrate (Oxidizing Agents)**

Emergency contact details
Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Registered, Manufactured and Distributed by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Waltloo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Spill Response and Transport incidents
Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Rolfes 
Agri

**Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use**



LIFT HERE

PRECAUTIONS:

- If medical advice is needed, have container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Obtain, read and follow all safety instructions before use.
- Keep away from heat, hot surfaces, sparks, open flames, and other ignition sources. No smoking.
- Keep away from clothing and other combustible product.
- Wear eye protection/ face protection.
- If exposed or concerned, get medical aid.
- In case of fire: Use suitable medium to extinguish
- Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler. Do not use empty containers for any other purpose.

Relevant Substances:

Boric Acid < 10% and Potassium Nitrate 30 - 60%

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

- **EASYGRO FLOWER & FRUIT** is recommended for use during plant growth periods that require relatively high potassium levels such as during flowering, fruit formation and fruit growth.
- **EASYGRO FLOWER & FRUIT** is a completely water-soluble fertilizer and chlorine free.
- **EASYGRO FLOWER & FRUIT** is readily taken up by the plant and is formulated for use as both a foliar feed and for fertigation.
- For optimum results **EASYGRO FLOWER & FRUIT** should be used in a program in combination with the other **EASYGRO range** of fertilizers.
- **EASYGRO CALCIUM** should be used in combination with **EASYGRO FLOWER & FRUIT** for increased fruit quality, shelf life and for correcting calcium and magnesium deficiencies.
- **EASYGRO FLOWER & FRUIT** is compatible with most agrochemicals. It is however advisable to do a miscibility test prior to mixing with other chemicals.

APPLICATION	RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Foliar feed	1 – 3 kg	1 – 2 kg	Vegetable and ornamental crops. Apply as a full cover spray. Apply at 1 – 2 week intervals throughout the flowering and fruiting period.
	1 – 3 kg	1 – 3 kg / 30 ℓ water	Aerial Application. Apply at 1 – 2 week intervals throughout the flowering and fruiting period.
	4 – 6 kg	1 – 2 kg	Large trees and orchards. Apply at 1 – 2 week intervals during flowering and fruiting until the last harvest.
Fertigation	5 – 10 kg	0,25 – 2 kg	Use lower concentrations young plants and increase the rate as the plants mature.

The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

EASYGRO FLOWER & FRUIT should preferably be applied in the early morning or late afternoon. **Do not apply to plants that are undergoing a period of moisture or heat stress.** Consult with your chemical distributor, consultant or manufacturer should crop specific programs or any other information be required around the use of the **EASYGRO range** of fertilizers.