

Easygro

Starter

Reg. No. K5241 Act No. 36 of 1947

FERTILIZER GROUP 1

READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

EasyGro Starter is a water soluble 2:1:2 (43) fertilizer with micro elements for foliar feeding and fertigation.



WARNING

Nitrogen (N)	175 g/kg	Cytokinins	0,5 mg/kg
Phosphorus (P)	85 g/kg	Auxins	1,0 mg/kg
Potassium (K)	174 g/kg		
Sulphur (S)	13 g/kg	*EDTA Chelated	
Magnesium (Mg)	3 g/kg		
*Zinc (Zn)	500 mg/kg		
*Iron (Fe)	500 mg/kg		
*Manganese (Mn)	200 mg/kg		
*Copper (Cu)	200 mg/kg		
Boron (B)	1000 mg/kg		
Molybdenum (Mo)	50 mg/kg		

HAZARD STATEMENTS:

May intensify fire; oxidizer.
May damage fertility or the unborn child.

PRECAUTIONARY STATEMENTS:

Keep away from heat, hot surfaces, sparks, open flames, and other ignition sources. No smoking.
If exposed or concerned, get medical aid.

Rolfes 
Agri

Registered, Manufactured and Distributed by: Rolfes Agri (Pty) Ltd.
(Reg. No. 1998/013411/07) • 288 Mundt Street, Walfloo, 0184
Gauteng, RSA. • Tel: (012) 803-0145

Emergency contact details

Office Hour Poisoning Helpline
Rolfes Agri (Pty) Ltd.
Tel: +27 (12) 803 0145

Spill Response and Transport incidents

Spill Tech, Oil and chemical pollution control
Tel: +27 (86) 100 0366 / +27 (83) 253 6618
www.spilltech.co.za

Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

Precautionary Statement:

- If medical advice is needed, have container or label at hand.
- Keep out of reach of children.
- Read carefully and follow all instructions.
- Keep away from heat, hot surfaces, sparks, open flames, and other ignition sources.
- No smoking.
- Obtain, read and follow all safety instructions before use.
- Keep away from clothing and other combustible product.
- Wear protective gloves/ Protective clothing/ Eye protection when handling this product.
- In case of fire: Use suitable medium to extinguish
- If exposed or concerned, get medical aid.
- Store locked up.
- Empty all contents from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds. Puncture the rinsed container to render it useless and send to a recycler. Do not use empty containers for any other purpose.

Relevant Substances:

Potassium Nitrate 30 – 60% and Boric Acid <10%.

EASYGRO STARTER is recommended for use during plant growth periods that require relatively high phosphorus levels such as seedlings, transplants, young plants and during periods of stress when active root growth is essential or has been reduced.

EASYGRO STARTER is a completely water-soluble fertilizer and chlorine free.

EASYGRO STARTER is readily taken up by the plant and is formulated for use as both a foliar feed and for fertigation.

For optimum results **EASYGRO STARTER** should be used in a program in combination with the other EASYGRO range of fertilizers.

In the case of Hydroponics, **EASYGRO STARTER** should be used in combination with **EASYGRO CALMAG + TE***.

EASYGRO STARTER is compatible with most agrochemicals. It is **however advisable to do a miscibility test prior to mixing with other chemicals.**

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

APPLICATION	RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Soil drench	3 – 6 kg	1 – 2 kg	Apply directly onto the soil and leach in with irrigation before planting or directly thereafter. Apply directly into planting hole as a root drench before transplanting.
Foliar Feed	1 – 3 kg	1 – 2 kg	Vegetable and ornamental crops. Apply as a full cover spray. Apply at 1 -2 week intervals for the first 4 weeks after establishment.
	1 – 3 kg	1 – 3 kg / 30 ℓ water	Aerial Application. Apply at 1 – 2 week intervals as required.
	4 – 6 kg	1 – 2 kg	Large trees and orchards. Apply at 1 – 2 week intervals prior to flowering and after last harvest.
Fertigation	5 – 10 kg	0,25 – 2 kg	Use lower concentrations for young plants and increase the rate as the plants increase in size. Use with EASYGRO Calcium in a program.

The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

EASYGRO STARTER should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress. Consult with your chemical distributor, consultant or manufacturer should crop specific programs or any other information be required around the use of the **EASYGRO** range of fertilizers.